

**TRANSACTIONS OF SOCIETY OF ACTUARIES
1970 REPORTS**

**III. AGE-ADJUSTED DEATH RATES FOR LEADING
CAUSES OF DEATH IN THE UNITED STATES**

TABLES 1 and 2 on the following pages show death rates per 100,000 from selected causes of death by year for the years 1950 through 1967 and update the tables shown on pages D461-D464 of Volume XIX of the *Transactions*. The rates are for all ages combined and are shown separately by sex for white persons in the United States (Table 1) and for nonwhite persons in the United States (Table 2). All rates are age-adjusted on the basis of the age distribution of the total population of the United States in the census of April 1, 1940.

The Committee on Mortality under Ordinary Insurances and Annuities is indebted to the Division of Vital Statistics, National Center for Health Statistics of the United States Department of Health, Education, and Welfare for having provided these data.

TABLE 1

DEATH RATES PER 100,000 AT ALL AGES* FROM SELECTED CAUSES
FOR WHITE PERSONS BY SEX, UNITED STATES, 1950-67

YEAR	MAJOR CARDIOVASCULAR-RENAL DISEASES						MALIGNANT NEOPLASMS	
	Total	Diseases of Cardiovascular System				Chronic Nephritis, etc.		
		Total	Vascular Lesions	Diseases of Heart	General Arteriosclerosis			
Males								
1967	469.0	464.7	71.5	361.6	12.3	4.3	150.3	
1966	480.2	475.6	73.6	369.8	13.2	4.6	148.9	
1965	479.3	474.5	73.8	368.6	13.4	4.8	148.0	
1964	479.2	474.1	74.6	367.7	13.1	5.0	145.6	
1963†	490.5	485.2	77.6	375.7	13.8	5.2	143.8	
1962†	484.5	479.1	77.4	370.3	13.7	5.5	142.0	
1961	480.5	474.9	77.3	367.3	13.7	5.7	142.0	
1960	493.2	487.3	80.3	375.4	14.7	5.9	141.6	
1959	487.3	481.2	80.9	369.4	14.4	6.0	139.4	
1958	495.1	488.2	82.9	373.7	14.9	6.9	138.2	
1957	496.8	489.2	84.4	375.6	14.8	7.6	139.9	
1956	488.8	481.1	82.1	370.2	14.7	7.8	139.1	
1955	487.0	478.6	82.7	367.4	15.6	8.4	137.4	
1954	480.4	471.3	82.0	362.2	14.8	9.0	136.0	
1953	501.5	491.3	85.6	376.5	16.7	10.2	135.5	
1952	501.9	490.4	86.4	374.7	16.8	11.5	133.7	
1951	508.3	495.7	87.3	378.0	17.9	12.6	131.1	
1950	514.1	499.9	87.0	381.1	18.4	14.2	130.9	
Females								
1967	263.0	260.3	59.9	179.2	10.3	2.6	107.9	
1966	272.0	269.1	62.0	185.2	10.9	2.9	107.6	
1965	272.9	269.8	62.1	185.9	10.9	3.0	108.1	
1964	270.1	272.9	63.3	187.9	11.1	3.3	107.6	
1963†	283.7	280.3	65.9	192.3	11.4	3.4	107.3	
1962†	283.3	279.8	66.5	191.6	11.5	3.5	107.4	
1961	281.6	277.8	66.3	190.5	11.2	3.7	108.5	
1960	291.5	287.6	68.7	197.1	11.7	3.9	109.5	
1959	292.6	288.5	70.2	196.6	11.7	4.1	109.4	
1958	301.6	296.9	72.5	202.4	11.9	4.8	110.6	
1957	305.3	299.8	73.4	205.8	11.9	5.5	112.6	
1956	301.9	296.2	72.3	203.5	11.9	5.7	113.2	
1955	303.8	297.8	73.2	204.0	12.4	6.0	114.3	
1954	303.2	296.4	73.5	202.6	12.1	6.9	115.1	
1953	321.5	313.8	77.7	214.1	13.4	7.7	116.4	
1952	325.1	316.7	78.4	215.7	13.9	8.4	117.6	
1951	349.8	339.7	83.9	218.9	16.5	10.0	119.1	
1950	339.0	328.0	79.7	223.6	14.3	11.0	119.4	

* Age-adjusted on the basis of the age distribution of the total population of the United States in the census of April 1, 1940.

† United States, excluding New Jersey.

TABLE 1—*Continued*
DEATH RATES PER 100,000 AT ALL AGES* FROM SELECTED CAUSES
FOR WHITE PERSONS BY SEX, UNITED STATES, 1950-67

YEAR	ACCIDENTS			INFLUENZA AND PNEU- MONIA	DIABETES MELLITUS	CIRRHOSIS OF THE LIVER	TUBERCU- LOSIS, ALL FORMS
	Total	Motor Vehicle	Other				
Males							
1967 . . .	77.8	41.1	36.7	24.2	12.4	17.2	3.6
1966 . . .	78.5	41.9	36.6	27.1	12.3	16.8	4.2
1965 . . .	75.4	39.4	36.0	26.9	11.9	15.6	4.5
1964 . . .	73.4	38.1	35.3	25.9	11.8	15.0	4.8
1963† . . .	72.4	36.7	35.6	30.7	11.9	14.7	5.4
1962† . . .	70.7	34.9	35.7	26.5	11.8	14.6	5.6
1961 . . .	68.5	33.4	35.2	24.8	11.4	14.2	6.1
1960 . . .	70.6	34.0	36.5	31.0	11.6	14.4	6.8
1959 . . .	71.2	34.5	36.6	25.8	11.0	13.7	7.3
1958 . . .	71.1	34.3	36.7	27.7	11.1	13.7	8.0
1957 . . .	76.1	36.8	39.2	30.5	10.9	14.4	8.9
1956 . . .	77.8	38.5	39.4	23.5	10.9	13.4	9.3
1955 . . .	77.7	37.8	40.0	22.3	10.9	12.9	10.0
1954 . . .	75.9	35.4	40.6	21.2	10.9	12.8	11.1
1953 . . .	81.8	38.0	43.8	27.5	11.2	13.0	13.2
1952 . . .	83.2	38.4	44.8	24.8	11.5	13.0	16.6
1951 . . .	84.5	37.8	46.7	26.6	11.2	12.3	21.1
1950 . . .	81.0	35.9	45.0	27.1	11.3	11.6	23.3
Females							
1967 . . .	28.2	14.4	13.8	14.3	12.8	8.4	1.1
1966 . . .	28.7	14.7	14.0	16.4	13.0	7.9	1.2
1965 . . .	27.7	13.7	14.0	16.1	12.9	7.6	1.3
1964 . . .	27.3	13.6	13.7	15.7	12.9	7.3	1.4
1963† . . .	26.7	12.6	14.1	19.0	13.3	7.2	1.6
1962† . . .	26.2	12.0	14.2	16.7	13.1	6.8	1.8
1961 . . .	24.5	10.8	13.6	15.5	13.3	6.7	1.9
1960 . . .	25.4	11.1	14.4	19.0	13.7	6.6	2.2
1959 . . .	25.1	11.0	14.1	16.2	13.5	6.4	2.4
1958 . . .	25.1	10.8	14.3	17.2	13.6	6.3	2.6
1957 . . .	27.0	11.2	15.8	19.3	14.3	6.6	2.9
1956 . . .	27.4	11.5	15.8	15.1	14.0	6.5	3.2
1955 . . .	27.6	11.4	16.2	14.9	14.1	6.1	3.6
1954 . . .	27.1	10.6	16.6	13.9	14.5	6.0	4.1
1953 . . .	29.1	11.5	17.6	18.2	15.6	6.4	5.0
1952 . . .	30.6	11.6	19.0	16.9	15.8	6.2	6.8
1951 . . .	32.0	11.2	20.8	19.7	16.6	6.2	9.2
1950 . . .	30.6	10.6	20.0	18.9	16.4	5.8	10.2

* Age-adjusted on the basis of the age distribution of the total population of the United States in the census of April 1, 1940.

† United States, excluding New Jersey.

TABLE 2

DEATH RATES PER 100,000 AT ALL AGES* FROM SELECTED CAUSES
FOR NONWHITE PERSONS BY SEX, UNITED STATES, 1950-67

YEAR	MAJOR CARDIOVASCULAR-RENAL DISEASES					MALIGNANT NEO-PLASMS	
	Total	Diseases of Cardiovascular System					
		Total	Vascular Lesions	Diseases of Heart	General Arteriosclerosis	Chronic Nephritis, etc.	
Males							
1967 . . .	551.0	537.8	124.4	370.7	11.8	13.2	186.6
1966 . . .	575.6	561.5	131.5	385.5	13.2	14.1	182.3
1965 . . .	569.8	555.8	134.2	376.1	13.6	14.0	173.3
1964 . . .	565.3	550.8	131.1	375.2	13.6	14.5	170.1
1963† . . .	584.7	569.6	135.7	387.0	14.7	15.2	168.7
1962† . . .	569.4	554.7	133.4	376.4	15.0	14.7	159.1
1961 . . .	548.3	533.5	127.5	362.7	14.2	14.8	157.9
1960 . . .	564.0	547.8	135.2	368.3	15.5	16.1	154.8
1959 . . .	562.2	545.4	134.8	365.9	15.2	16.8	152.5
1958 . . .	582.1	563.4	141.2	377.1	16.0	18.7	146.4
1957 . . .	589.7	569.8	140.2	387.2	15.2	20.0	149.2
1956 . . .	568.2	547.2	134.3	372.7	14.8	21.0	145.0
1955 . . .	567.3	546.7	136.2	369.2	16.2	20.5	138.7
1954 . . .	567.4	542.7	136.0	367.0	15.5	24.7	140.6
1953 . . .	611.2	584.6	141.7	398.3	17.7	26.6	133.2
1952 . . .	621.4	588.6	143.9	402.8	17.1	32.9	130.7
1951 . . .	619.9	586.0	142.6	400.5	17.1	33.9	125.3
1950 . . .	633.7	595.4	144.0	407.5	18.0	38.3	125.8
Females							
1967 . . .	417.3	406.7	114.1	257.8	9.7	10.7	126.6
1966 . . .	437.8	427.4	120.3	270.5	11.4	10.4	127.0
1965 . . .	440.8	430.1	125.5	269.2	11.3	10.7	125.2
1964 . . .	450.2	438.5	125.8	276.4	11.8	11.6	124.0
1963† . . .	466.8	455.3	131.7	286.9	12.4	11.5	124.4
1962† . . .	463.2	451.2	132.2	283.0	12.1	12.0	124.7
1961 . . .	455.0	443.4	130.2	278.0	11.5	11.7	124.3
1960 . . .	467.1	453.8	134.4	283.3	12.3	13.2	125.0
1959 . . .	465.4	451.2	136.5	279.2	12.3	14.2	121.3
1958 . . .	492.4	476.2	139.4	297.5	13.7	16.3	125.3
1957 . . .	503.6	485.2	142.2	307.6	13.1	18.4	125.2
1956 . . .	490.8	471.8	139.1	298.7	12.5	19.0	127.8
1955 . . .	486.1	466.5	139.3	293.0	12.5	19.6	124.7
1954 . . .	489.4	469.0	142.0	292.1	12.6	20.4	125.9
1953 . . .	520.9	496.2	145.7	314.6	13.5	24.7	125.1
1952 . . .	538.6	509.6	149.9	323.0	13.1	29.1	126.8
1951 . . .	550.2	518.6	152.1	329.5	13.1	31.6	126.4
1950 . . .	570.8	534.7	153.4	342.9	13.7	36.1	131.0

* Age-adjusted on the basis of the age distribution of the total population of the United States in the census of April 1, 1940.

† United States, excluding New Jersey.

TABLE 2—Continued

DEATH RATES PER 100,000 AT ALL AGES* FROM SELECTED CAUSES
FOR NONWHITE PERSONS BY SEX, UNITED STATES, 1950-67

YEAR	ACCIDENTS			INFLUENZA AND PNEU- MONIA	DIABETES MELLITUS	CIRRHOSIS OF THE LIVER	TUBERCU- LOSIS, ALL FORMS
	Total	Motor Vehicle	Other				
Males							
1967....	114.7	49.4	65.3	47.6	18.5	27.2	15.4
1966....	116.4	50.1	66.2	55.2	18.3	26.6	16.3
1965....	109.3	46.4	63.0	52.3	18.1	23.3	16.0
1964....	105.2	44.2	61.0	51.2	17.6	19.6	17.0
1963†....	103.4	42.7	60.8	66.6	16.6	17.1	18.8
1962†....	100.5	40.3	60.2	52.1	16.1	16.8	19.2
1961....	95.1	38.0	57.1	49.6	14.9	15.9	19.9
1960....	101.1	39.5	61.6	68.0	16.1	14.9	21.4
1959....	100.2	40.5	59.7	53.6	14.1	13.7	22.6
1958....	100.2	40.4	59.7	61.4	13.0	12.5	26.5
1957....	106.5	43.5	63.0	65.0	12.5	13.4	27.6
1956....	109.7	47.9	61.7	49.1	11.7	11.8	29.5
1955....	106.5	45.1	61.5	50.8	11.2	10.3	32.1
1954....	104.6	42.6	62.0	49.1	11.3	9.8	35.8
1953....	116.2	47.8	68.4	68.6	11.3	10.0	45.8
1952....	118.2	46.9	71.4	60.6	11.5	10.6	59.6
1951....	114.1	45.2	68.8	62.3	11.8	9.9	73.6
1950....	107.1	41.2	66.0	63.4	11.8	9.0	81.8
Females							
1967....	35.9	14.6	21.3	26.6	29.7	15.0	5.5
1966....	39.1	15.0	24.1	32.8	30.5	15.0	6.0
1965....	35.8	13.7	22.1	32.3	28.6	13.4	6.4
1964....	34.9	12.5	22.5	31.4	29.0	11.3	6.7
1963†....	36.1	11.8	24.3	45.1	29.1	11.0	7.3
1962†....	35.4	11.4	24.0	35.4	27.1	11.0	8.1
1961....	33.9	11.0	23.0	30.5	26.7	9.9	8.5
1960....	36.1	10.6	25.5	43.3	26.8	9.1	9.3
1959....	34.6	10.8	23.8	35.2	24.2	9.1	10.9
1958....	35.8	10.2	25.6	39.0	24.3	7.4	11.9
1957....	38.4	12.5	25.9	46.4	23.6	8.4	12.8
1956....	37.3	12.5	24.8	34.6	22.2	7.8	14.8
1955....	38.5	12.6	25.8	33.9	21.6	6.8	16.5
1954....	37.8	11.8	26.0	33.8	21.2	6.5	19.1
1953....	39.8	12.3	27.5	49.1	22.9	6.3	23.9
1952....	41.6	12.4	29.2	44.0	21.5	5.7	32.7
1951....	41.0	12.5	28.6	47.2	21.3	6.3	42.8
1950....	38.8	11.1	27.7	50.6	22.6	5.9	53.2

* Age-adjusted on the basis of the age distribution of the total population of the United States in the census of April 1, 1940.

† United States, excluding New Jersey.