



SOCIETY OF ACTUARIES

Article from:

Long-Term Care News

September 2008 – Issue No. 21

Is the Compression of Morbidity a Universal Phenomenon?

by Jean-Marie Robine, Siu Lan K. Cheung, Shiro Horiuchi and A. Roger Thatcher

Jean-Marie Robine
INSERM,
Health and Demography
Correspondence:
robine@valdorel.fnclcc.fr

Siu Lan K. Cheung
Hong Kong University of
Science and Technology,

Shiro Horiuchi
Program in Urban Public
Health at Hunter College
and CUNY Institute for
Demographic Research

A. Roger Thatcher
Former director of the
Office of Population Censuses
and Surveys, London

Acknowledgment: This study of the compression of morbidity is part of a wider project called the M-project, led by the four authors, which aims to explore the demographic factors leading to an increase in the adult longevity, investigate the phenomenon called the compression of mortality and develop a new methodological approach to longevity analysis with focus on the modal age at death (M).

ABSTRACT:

Recent reviews of national health trends show conflicting results. It is clear today that the various health dimensions follow different trends over time. For instance, an expansion of morbidity may accompany a compression of disability. What do we measure when we observe a decline in disability? Is the elderly population intrinsically healthier or are individuals more independent and less helped by children, using more technical devices in a more favorable environment? The recent OECD study, reviewing trends in ADL disability at age 65 and over in 12 OECD countries during the 1990s, demonstrates that there is clear evidence of a decline in disability among elderly people in only five of the 12 countries studied: Denmark, Finland, Italy, the

Netherlands and the United States. Three countries (Belgium, Japan and Sweden) report an increasing rate and two countries (Australia, Canada) a stable rate. In France and the United Kingdom, different surveys show different trends in ADL disability (OECD, 2007). These results suggest that a decline in ADL disability may be less universal than expected. More importantly, the OECD study shows that ADL disability at age 65 and over ranges widely from a low 7.1 percent in the Netherlands (HIS) to a high of 18 percent in the United Kingdom (GHS survey). This paper reviews available evidence about the compression of morbidity and the disability decline and discusses the context in which they occur: initial level of disability, initial value of life expectancy and trend in life expectancy. ■

