STABILITY OF PENSION SYSTEMS WHEN RATES OF RETURN ARE RANDOM

Daniel DUFRESNE

Abstract

Consider a funded pension plan, and suppose actuarial gains or losses are amortized over a fixed number of years. The paper aims at assessing how contributions (C) and fund levels (F) are affected when the rates of return of the plans's assets form an i.i.d. sequence of random variables. This is achieved by calculating the mean and variance of C_t and F_t for $t \leq \infty$.

Abbreviated title: PENSION SYSTEMS

Reywords: Pension funding, Random rates of return, Actuarial gains and losses.

- 81 -

1. Introduction

This paper examines the effect of random rates of return on pension fund levels and contributions. The funding methods considered are those which

- determine an actuarial liability (AL) and a normal cost (NC) at every valuation date; and
- (2) amortize individual inter-valuation gains or losses over a fixed number of years (e.g. 5 ou 15).

These methods have been used by actuaries in Canada and the United States.

<u>Remarks</u> 1. A similar study has been done of funding methods which satisfy (1) but adjust the normal cost by a constant fraction of the actuarial liability. See Dufresne (1986a) and (1988).

2. Pension mathematics and gain and loss analysis are described in Trowbridge (1952), Winklevoss (1977) and Lynch (1979).

- 82 -

•

2. Notation

AL	Actuarial liability
В	Benefit outgo
с	Overall contribution
F	Fund level
i	Valuation rate of interest
Lt	Actuarial loss during (t-1,t)
m	Amortization period for actuarial losses
NC	Normal cost
Р	Adjustment of normal cost
r	Mean rate of return on assets
rt	Rate of return on assets during (t-1,t)
ULt	Unfunded liability at time t
UL_{i}^{A}	Unfunded liability at time t if all actuarial assumptions work out during (t-1,t).
β _k	$= a_{m-k} / \ddot{a}_{m} , 1 \le k \le m-1$
λ _k	= ä /ä , 0 ≤ k ≤ m−1
σ²	Variance of r _t

3. Description of model and basic equations

In order to isolate the effect of fluctuating rates of return (and keep the model tractable), the following assuptions are made.

- I. The population is stationary form the start.
- II. Except for rates of return, all actuarial assumptions are consistently realized.
- III. There is no inflation on benefits.
- IV. The rates of return $\{r_t, t\}$ form an i.i.d. sequence with mean r and variance σ^2 . r_t is the rate earned on assets during the period (t-1,t).

Suppose the pension plan is set up at time t = 0. Given the assumptions above, the assets process satisfies

$$F_{t} = (1+r_{t})(F_{t-1}+C_{t-1}-B), t \ge 1$$
 (1)

where F is the fund level, C the contribution and B the benefit outgo. B is constant from assumptions I to III. On the liabilities side we have

$$AL = (1+i)(AL+NC-B)$$
(2)

where i is the valuation rate of interest. This equation is known as the equation of equilibrium.

Now define the unfunded liability at time t as $UL_t = AL-F_t$, t > 0, and the (actuarial) loss experienced during the period (t-1,t) as

 $L_t \approx UL_t - [value of UL_t had all actuarial assumptions been realized during (t-1,t)].$

$$= UL_{t} - UL_{t}^{n}, t \ge 1$$
(3)

For the time being let $L_t = 0$ for $t \le 0$. Letting $r_t = i$ in Eq. (1), and subtracting it from Eq. (2), we get

- 84 -

$$UL_{t}^{A} = (1+i)(UL_{t-1}+NC-C_{t-1}).$$
(4)

Under the funding methods considered, the contribution at time

t is

$$C_t = NC + P_t, \tag{5}$$

$$P_{t} = \sum_{k=0}^{m-1} L_{t-k} / \ddot{a}_{m} .$$
(6)

Here m (an integer) is the amortization period and

$$\ddot{\mathbf{a}}_{-} = [1 - (1 + i)^{-m}] / [1 - (1 + i)^{-1}].$$

Each L_s is thus liquidated by m payments of amount L_s/\ddot{a}_m , made at valuation dates s, s+1,...,s+m-1. The fact that \ddot{a}_m is calculated at rate i ensures that L_s is in fact cancelled out after the m-th payment is made.

<u>Remark</u>. It should be observed that all losses are assumed to be amortized in the same fashion, irrespective of their sign. In practice, it may happen that gains (i.e. negative losses) be written off immediately, in order to reduce the unfunded liability or the current contribution.

As they stand, the above equations do not permit the calculation of the moments of F and C. One way to proceed is as follows: (i) derive a difference equation involving the L's only;

(ii) calculate the moments of the L's from this equation;

(iii) finally, obtain the moments of F and C from those of the L's.

First, let us express $\ensuremath{\text{UL}_t}$ in terms of the L's. We have

$$UL_{t} = AL-F_{t}$$

$$= (1+i)(AL+NC-B)$$

$$- (1+r_{t})(F_{t-1}+NC+P_{t-1}-B)$$

$$= (1+r_{t})(AL-F_{t-1}-P_{t-1})$$

$$- (r_{t}-i)(AL+NC-B)$$

$$= (1+i)(UL_{t-1}-P_{t-1})$$

$$+ (r_{t}-i)(UL_{t-1}-P_{t-1}-(1+i)^{-1}AL)$$
(7)

In view of Eqs. (3), (4) and (5), this implies

$$L_{t} = (r_{t}-i)(UL_{t-1}-P_{t-1}-(1+i)^{-1}AL), \quad t \ge 1$$
(8)

Eq. (7) can be rewritten as

$$UL_{t} = (1+i)(UL_{t-1}-P_{t-1}) + L_{t}$$
(9)

or

$$UL_{t} - (1+i)UL_{t-1} = L_{t} - (1+i)\sum_{\substack{s=t-m \\ s=t-m}}^{t-1} L_{s} / \ddot{a}_{m}.$$
 (10)

A particular solution of this difference equation is

$$UL_{t}^{p} = \sum_{k \ge 0} \lambda_{k} L_{t-k}$$

where the λ 's can be determined by direct substitution into Eq. (10), yielding

$$\lambda_{0}L_{t} + [\lambda_{1} - (1+i)\lambda_{0}]L_{t-1} + [\lambda_{2} - (1+i)\lambda_{1}]L_{t-2} + \dots$$
$$= L_{t} - [(1+i)/\ddot{a}_{m}]L_{t-1} - \dots - [(1+i)/\ddot{a}_{m}]L_{t-m}$$

which means

- 86 -

)

1

ł

 $\lambda_{0} = 1$ $\lambda_{1} = \ddot{a}_{m-1} / \ddot{a}_{m}$ $\lambda_{2} = \ddot{a}_{m-2} / \ddot{a}_{m}$ \vdots $\lambda_{m-1} = \ddot{a}_{1} / \ddot{a}_{m}$ $\lambda_{k} = 0, k \ge m.$

A solution of the homogeneous equation

$$UL_{t} - (1+i)UL_{t-1} = 0$$

is $c(1+i)^t$, c a constant. The solution of the complete equation (10) is therefore (Brand (1966), p. 368)

$$UL_{t} = \sum_{k=0}^{m-1} \lambda_{k} L_{t-k} + UL_{o}(1+i)^{t}.$$

The term $UL_o(1+i)^{t}$ brings to light the fact that the initial unfunded liability ($UL_o = AL-F_o$) has not been taken into account so far. It is easy to see that including supplementary payments of amount $UL_o/\ddot{a_n}$ at times 0, 1,...,n-1 will liquidate UL_o entirely. For the sake of simplicity, let us assume that n = m, so we can define $L_o = UL_o$ and obtain

$$UL_{t} = \sum_{k=0}^{m-1} \lambda_{k} L_{t-k}, \quad t \ge 0$$
(11)

Eqs. (6), (8) and (11) now permit the derivation of a difference equation involving the L's only:

$$L_{t} = (r_{t}-i) \left[\sum_{k=0}^{m-1} (\lambda_{k}-1/\ddot{a}_{m}) L_{t-1-k} - (1+i)^{-1} AL \right]$$

= $(r_{t}-i) \left[\sum_{k=1}^{m-1} \beta_{k} L_{t-k} - (1+i)^{-1} AL \right]$ (12)

where $\beta_k = \lambda_{k-1} - 1/\ddot{a}_m = a_{m-k}/\ddot{a}_m$ (clearly $\beta_m = \lambda_{m-1} - 1/\ddot{a}_m = 0$).

- 87 -

4. Stability conditions

Definition. A sequence {y_t} satisfying

$$y_t + \sum_{j=1}^{n} \alpha_j y_{t-j} = w, t \ge 1$$
(13)

will be said to be <u>stable</u> if there is a finite value y^* such that $y_t \rightarrow y^*$ as $t \rightarrow \infty$ for any set of initial conditions $y_0, y_{-1}, \dots, y_{-n+1}$.

It is well known that a necessary and sufficient condition for this kind of stability is that all the roots of the characteristic equation

$$p(z) = z^{n} + \sum_{j=1}^{n} \alpha_{j} z^{n-j} = 0$$

be less than one in modulus.

<u>Proposition 1.</u> If $\sum |\alpha_j| < 1$, then $\{y_t\}$ is stable.

<u>Proof</u>. Suppose there exists $z \in C$ such that p(z) = 0 and $|z| \ge 1$. Then

$$\begin{array}{cccc} n & n & n \\ |z|^n & \sum |\alpha_j| & |z|^{n-j} & |z|^n & \sum |\alpha_j| & \langle |z|^n \\ j=1 & j=1 \end{array}$$

a contradiction. D

<u>Proposition 2</u>. Suppose $\alpha_j \leq 0$ for all j. Then $\{y_t\}$ is stable if and only if $|\sum \alpha_j| \leq 1$.

<u>Proof</u>. Sufficiency is a consequence of Prop. 1. To prove necessity, suppose $|\sum_{\alpha_i}| \ge 1$, and let $q(z) = z^n p(1/z)$. Then q(0) = 1 and

$$q(1) = 1 + \sum \alpha_j \leq 0.$$

- 88 -

ì

Thus q(z) has at least one root z^* in (0,1], which implies that p(z) has the root $z^{**} = 1/z^*$ in $[1, \infty)$.

<u>Remark</u>. That $|\sum \alpha_j| \le 1$ is not in general a necessary condition for stability of (13) can be seen by considering the case i = 0, m = 3 and $r_t - i \equiv p$ in Eq. (12),

$$L_t - (2p/3)L_{t-1} - (p/3)L_{t-2} = pAL.$$

This sequence is stable for $-3 , while <math>|\sum \alpha_j| = |p|$.

Let us now return to the processes
$$\{F_t\}$$
 and $\{C_t\}$.

<u>Definition</u>. A process $\{X_t\}$ will be said to be <u>p-th order stable</u> if $\{EX_t^p\}$ is stable.

Since

$$F_t = AL - UL_t$$

$$= AL - \sum_{k=0}^{m-1} \lambda_k L_{t-k},$$

$$C_{t} \approx NC + \sum_{k=0}^{m-1} L_{t-k} / \ddot{a}_{m},$$

it is evident that the stability properties of $\{F_t\}$ and $\{C_t\}$ are the same as those of $\{L_t\}$. We will thus consider Eq. (12), with initial conditions being now arbitrary (imagining that the plan has been in existence for some time before t = 0).

- 89 -

We get

$$EL_{t} = E(r_{t}-i) \left\{ \sum_{k=1}^{m-1} \beta_{k} EL_{t-k} - (1+i)^{-1}AL \right\}$$

since r_t is independent of L_{t-k} , $k \ge 1$. Applying Prop. 1, we obtain

<u>Proposition 3</u>. If $|r-i|\Sigma \beta_k < 1$, then $\{L_t\}$, $\{F_t\}$ and $\{C_t\}$ are first order stable, and

- (a) $\lim_{t\to\infty} EL_t = M_{\infty} = -(r-i)(1+i)^{-1}AL/(1-(r-i)\Sigma\beta_k),$
- (b) $\lim_{t\to\infty} EF_t = AL M_{\infty} \sum \lambda_k$,
- (c) $\lim_{t\to\infty} EC_t = NC + M_{\infty} m/\ddot{a}_m$.

Second order stability

At this point we make a supplementary assumption:

V. $Er_t = r = i$.

Using Eq. (12), this implies

$$EL_{+} = 0, t \ge 1$$

and

$$EL_{t}L_{s} = E(r_{t}-i) E\left(\sum_{k=1}^{m-1} \beta_{k} L_{t-k} - (1+i)^{-1} AL\right) L_{s}$$
$$= 0$$

for any $t \ge 1$, s < t. Thus $\{L_t, t \ge 1\}$ becomes a sequence of uncorrelated zero-mean r.v.'s. Eq. (12) then gives

- 90 -

$$\operatorname{Var} L_{t} = \operatorname{EL}_{t}^{2} = \sigma^{2} \left(\sum_{k=1}^{m-1} \beta_{k}^{2} \operatorname{Var} L_{t-k} + (1+i)^{-2} \operatorname{AL}^{2} \right), t \ge 1$$

<u>Proposition 4</u>. {L_t}, {F_t} and {C_t} are second order stable if and only if $\sigma^2 \sum \beta_k^2 < 1$, in which case

$$\begin{split} &\lim_{t\to\infty} \operatorname{Var} L_t = \operatorname{V}_{\mathfrak{W}} = \sigma^2 (1+i)^{-2} \operatorname{AL}^2 / (1-\sigma^2 \sum \beta_k^2) \\ &\lim_{t\to\infty} \operatorname{Var} F_t = \operatorname{V}_{\mathfrak{W}} \sum \lambda_k^2 \\ &\lim_{t\to\infty} \operatorname{Var} C_t = \operatorname{V}_{\mathfrak{W}} m / (\ddot{a}_{\overline{m}})^2 . \end{split}$$

<u>Remarks</u> 1. The L's are uncorrelated but certainly not independent. For example, let m = 2, r = i = 0, AL = 1/2. Then $\beta_1 = 1/2$ and

 $L_{t} = x_{t}(L_{t-1}-1)$

where $x_t = r_t/2$. If furthermore L(0) = 0 and $P(x_t = x) = P(x_t = -x) = 1/2$, we get

$$P(L_{t} = -x - x^{2} - ... - x^{t}) = P(x_{s} = x, 1 \le s \le t)$$
$$= (1/2)^{t},$$
$$P(L_{1} = x) = 1/2$$

and

1

ł

- 91 -

$$P(L_1 = x, L_1 = -x - x^2 - ... - x^t) = 0.$$

2. Covariances may also be calculated, yielding

$$Cov(F_{t}, F_{t+h}) = \sum_{k=0}^{m-h-1} Var(L_{t-k})\ddot{a}_{m-k} \ \ddot{a}_{m-k-h} / (\ddot{a}_{m})^{2}, \ 0 \le h \le m$$
$$= 0, \qquad h \ge m$$
$$Cov(C_{t}, C_{t+h}) = \sum_{k=0}^{m-h-1} Var(L_{t-k}) / (\ddot{a}_{m})^{2}, \ 0 \le h \le m$$
$$= 0, \qquad h \ge m.$$

5. Numerical illustration

The purpose of this section is to illustrate the results of Prop. 4.

Population	English Life Table No. 13 (males), stationary				
Entry age	30 (only)				
Retirement age	65				
No salary scale, no inflation on salaries					
Benefits	Straight life annity (2/3 of salary)				
Funding method	Entry Age Normal				
Valuation rate of interest	i = .01				
Actuarial liability	AL = 451% of payroll				
Normal cost	NC = 14.5% of payroll				
Earned rates of return	${r_t,t\geq 1}$ i.i.d. with $Er_t = r = .01$				

1

1

\$

I

- 92 -

Because $\text{Er}_t = i$, $\text{EF}_t = AL$ and $\text{EC}_t = NC$ for $t \ge m$, for any initial conditions. Table 1 contains the limiting coefficients of variation of F_t and C_t , that is to say

lim [Var
$$F_t$$
]^{1/2}/AL
t+ ∞
lim [Var C_t]^{1/2}/NC ,

. 1.

and

for various values of **m** and
$$\sigma = [Var r_t]^{1/2}$$
.

m	m = .025		σ = .05		<u>σ = .10</u>	
	1	1	1	1	1	1
	$\left[Var F(\omega) \right]^2$	$\left[\operatorname{Var} C(\boldsymbol{\omega})\right]^2$	$\left[Var F(m) \right]^{2}$	[Var C(m)] ²	$\left[\operatorname{Var} F(\mathfrak{o})\right]^2$	$[Var C(\omega)]^2$
	AL.	NC	AL	NC	AL	NC
1	2.5 %	77.0%	5.0 %	154.0 %	9.9 %	307.8 %
5	3.7	35.1	7.4	70.3	14.8	141.3
10	4.9	25.5	9.9	51.1	19.9	103.2
20	6.8	18.9	13.7	38.1	28.0	78.1
40	9.7	14.7	19.6	29.9	41.6	63.3

<u>TABLE 1</u>. Coefficients of variation of $F(\omega)$ and $C(\omega)$ (Er(t) = 0.01 , $\sigma = [Var r(t)]^{1/2}$)

Comments

1. It is seen that for $\sigma \leq 10$ %, the standard deviations of F_{00} and C_{00} are nearly linear in σ . This linearity gradually disappears, though, as σ or m become larger.

2. Within the range of σ and m chosen, no single value of m is "better" than the others. As m is varied, there is a trade-off between Var F and Var C, e.g. incereasing m reduces Var C, but increases Var F. 3. This trade-off is a direct outcome of Prop. 4. However, the following approximate formulas give a more intuitive understanding of the way Var F and Var C vary with m. They are valid when i = 0 and $\sigma^2 m$ is small (see proof below):

$$\operatorname{Var} F_{\omega} \approx \sigma^2 \frac{m}{3} AL^2 \tag{14}$$

$$\operatorname{Var} C_{to} \approx \sigma^2 \frac{1}{m} \operatorname{AL}^2$$
(15)

In words: when i is close to 0, the standard deviation of F (resp. of C) is roughly proportional to $m^{1/2}$ (resp. to $1/m^{1/2}$). For instance, in Table 1, moving from m = 5 to m = 20 approximately doubles st. dev. F_m and halves st.dev. C_m.

Proof of Eqs. (14) and (15). Set i = 0 in Prop. 4 to get

$$V_{00} = \sigma^2 AL^2 / \left(1 - \sigma^2 \sum_{k=1}^{m-1} \left[(m-k)/m \right]^2 \right)$$

$$Var F_{\omega} = V_{\omega} \sum_{k=0}^{m-1} [(m-k)/m]^2$$

Var
$$C_{\infty} = V_{\infty}/m$$

First,
$$m-1 = m-1$$

 $\sum_{k=1}^{m-1} [(m-k)/m]^2 = \sum_{j=1}^{m-1} j^2/m^2$

= $[(m-1)m(2m-1)/6]/m^2$

This shows that $V_{\infty} = \sigma^2 AL^2$ if $\sigma^2 m$ is small. Observing that similarly

$$\sum_{k=0}^{m-1} [(m-k)/m]^2 = m/3$$

1

we obtain Eqs. (14) and (15).

I.

1

;

,

;

1

,

1

1

<u>Remark</u>. As approximations for Var F_{∞} and Var C_{∞} , Eqs. (14) and (15) are sometimes valuable, even when i $\neq 0$. For example, if i = .01, σ = .05 and m = 10, Eq. (14) yields

$$[Var F_{co}]^{1/2}/AL = 9.1\%$$

while the exact number is 9.9% (Table 1).

ACKNOWLEDGMENTS

This paper is based on part of my Ph.D thesis. I wish to thank my supervisor, Prof. Steve Haberman, of the City University.

REFERENCES

Brand, L. (1966) Differential and Difference Equations. Wiley, New York.

- Dufresne, D. (1986a). Pension funding and random rates of return. In: Insurance and Risk Theory, Eds. M. Goovaerts et al., Riedel, Dordrecht.
- Dufresne, D. (1986b). The Dynamics of Pension Funding. Ph.D. thesis, The City University, London.
- Dufresne, D. (1985). Moments of pension fund contributions and fund levels when rates of return are random. To appear in <u>The Journal of the Institute</u> of Actuaries.
- Lynch, J.M. (1979). Identifying actuarial gains and losses in pension plans. Proceedings of the conference of Actuaries in Public Practice 26, 582-594.
- Trowbridge, C.L. (1952). Fundamentals of pension funding. <u>Transactions of the</u> Society of Actuaries 4, 17-43.
- Winklevoss, H.E. (1977) <u>Pension Mathematics</u>: <u>With Illustrations</u>. Irwin, Homewood, Illinois.