

SOCIETY OF ACTUARIES

EXAM M ACTUARIAL MODELS

EXAM M SAMPLE QUESTIONS

Copyright 2005 by the Society of Actuaries

Some of the questions in this study note are taken from past SOA examinations.

M-09-05

PRINTED IN U.S.A.

1. For two independent lives now age 30 and 34, you are given:

x	q_x
30	0.1
31	0.2
32	0.3
33	0.4
34	0.5
35	0.6
36	0.7
37	0.8

Calculate the probability that the last death of these two lives will occur during the 3rd year from now (i.e. ${}_2|q_{30:34}$).

- (A) 0.01
- (B) 0.03
- (C) 0.14
- (D) 0.18
- (E) 0.24

2. For a whole life insurance of 1000 on (x) with benefits payable at the moment of death:

(i)
$$\delta_t = \begin{cases} 0.04, & 0 < t \leq 10 \\ 0.05, & 10 < t \end{cases}$$

(ii)
$$\mu_x(t) = \begin{cases} 0.06, & 0 < t \leq 10 \\ 0.07, & 10 < t \end{cases}$$

Calculate the single benefit premium for this insurance.

(A) 379

(B) 411

(C) 444

(D) 519

(E) 594

3. A health plan implements an incentive to physicians to control hospitalization under which the physicians will be paid a bonus B equal to c times the amount by which total hospital claims are under 400 ($0 \leq c \leq 1$).

The effect the incentive plan will have on underlying hospital claims is modeled by assuming that the new total hospital claims will follow a two-parameter Pareto distribution with $\alpha = 2$ and $\theta = 300$.

$$E(B) = 100$$

Calculate c .

(A) 0.44

(B) 0.48

(C) 0.52

(D) 0.56

(E) 0.60

4. Computer maintenance costs for a department are modeled as follows:

- (i) The distribution of the number of maintenance calls each machine will need in a year is Poisson with mean 3.
- (ii) The cost for a maintenance call has mean 80 and standard deviation 200.
- (iii) The number of maintenance calls and the costs of the maintenance calls are all mutually independent.

The department must buy a maintenance contract to cover repairs if there is at least a 10% probability that aggregate maintenance costs in a given year will exceed 120% of the expected costs.

Using the normal approximation for the distribution of the aggregate maintenance costs, calculate the minimum number of computers needed to avoid purchasing a maintenance contract.

- (A) 80
- (B) 90
- (C) 100
- (D) 110
- (E) 120

5. A whole life policy provides that upon accidental death as a passenger on an airplane a benefit of 1,000,000 will be paid. If death occurs from other accidental causes, a death benefit of 500,000 will be paid. If death occurs from a cause other than an accident, a death benefit of 250,000 will be paid.

You are given:

- (i) Death benefits are payable at the moment of death.
- (ii) $\mu^{(1)} = 1/2,000,000$ where (1) indicates accidental death as a passenger on an airplane.
- (iii) $\mu^{(2)} = 1/250,000$ where (2) indicates death from other accidental causes.
- (iv) $\mu^{(3)} = 1/10,000$ where (3) indicates non-accidental death.
- (v) $\delta = 0.06$

Calculate the single benefit premium for this insurance.

- (A) 450
- (B) 460
- (C) 470
- (D) 480
- (E) 490

6. For a special fully discrete whole life insurance of 1000 on (40):

- (i) The level benefit premium for each of the first 20 years is π .
- (ii) The benefit premium payable thereafter at age x is $1000vq_x$, $x = 60, 61, 62, \dots$
- (iii) Mortality follows the Illustrative Life Table.
- (iv) $i = 0.06$

Calculate π .

- (A) 4.79
- (B) 5.11
- (C) 5.34
- (D) 5.75
- (E) 6.07

7. For an annuity payable semiannually, you are given:

- (i) Deaths are uniformly distributed over each year of age.
- (ii) $q_{69} = 0.03$
- (iii) $i = 0.06$
- (iv) $1000\bar{A}_{70} = 530$

Calculate $\ddot{a}_{69}^{(2)}$.

- (A) 8.35
- (B) 8.47
- (C) 8.59
- (D) 8.72
- (E) 8.85

8. For a sequence, $u(k)$ is defined by the following recursion formula

$$u(k) = \alpha(k) + \beta(k) \times u(k-1) \text{ for } k = 1, 2, 3, \dots$$

(i) $\alpha(k) = -\left(\frac{q_{k-1}}{p_{k-1}}\right)$

(ii) $\beta(k) = \frac{1+i}{p_{k-1}}$

(iii) $u(70) = 1.0$

Which of the following is equal to $u(40)$?

(A) A_{30}

(B) A_{40}

(C) $A_{40:\overline{30}}$

(D) $A_{40:\overline{30}}^1$

(E) $A_{40:\overline{30}}^{\frac{1}{i}}$

9. Subway trains arrive at a station at a Poisson rate of 20 per hour. 25% of the trains are express and 75% are local. The type of each train is independent of the types of preceding trains. An express gets you to the stop for work in 16 minutes and a local gets you there in 28 minutes. You always take the first train to arrive. Your co-worker always takes the first express. You both are waiting at the same station.

Calculate the probability that the train you take will arrive at the stop for work before the train your co-worker takes.

- (A) 0.28
- (B) 0.37
- (C) 0.50
- (D) 0.56
- (E) 0.75

10. For a fully discrete whole life insurance of 1000 on (40), the contract premium is the level annual benefit premium based on the mortality assumption at issue. At time 10, the actuary decides to increase the mortality rates for ages 50 and higher.

You are given:

- (i) $d = 0.05$
- (ii) Mortality assumptions:

At issue	${}_k q_{40} = 0.02, k = 0, 1, 2, \dots, 49$
Revised prospectively at time 10	${}_k q_{50} = 0.04, k = 0, 1, 2, \dots, 24$

- (iii) ${}_{10}L$ is the prospective loss random variable at time 10 using the contract premium.

Calculate $E[{}_{10}L | K(40) \geq 10]$ using the revised mortality assumption.

- (A) Less than 225
- (B) At least 225, but less than 250
- (C) At least 250, but less than 275
- (D) At least 275, but less than 300
- (E) At least 300

11. For a group of individuals all age x , of which 30% are smokers and 70% are non-smokers, you are given:

- (i) $\delta = 0.10$
- (ii) $\bar{A}_x^{\text{smoker}} = 0.444$
- (iii) $\bar{A}_x^{\text{non-smoker}} = 0.286$
- (iv) T is the future lifetime of (x) .
- (v) $\text{Var}\left[\bar{a}_{\overline{T}|}^{\text{smoker}}\right] = 8.818$
- (vi) $\text{Var}\left[\bar{a}_{\overline{T}|}^{\text{non-smoker}}\right] = 8.503$

Calculate $\text{Var}\left[\bar{a}_{\overline{T}|}\right]$ for an individual chosen at random from this group.

- (A) 8.5
- (B) 8.6
- (C) 8.8
- (D) 9.0
- (E) 9.1

12. T , the future lifetime of (0), has a spliced distribution.

- (i) $f_1(t)$ follows the Illustrative Life Table.
- (ii) $f_2(t)$ follows DeMoivre's law with $\omega = 100$.
- (iii) $f_T(t) = \begin{cases} k f_1(t), & 0 \leq t \leq 50 \\ 1.2 f_2(t), & 50 < t \end{cases}$

Calculate ${}_{10}P_{40}$.

- (A) 0.81
- (B) 0.85
- (C) 0.88
- (D) 0.92
- (E) 0.96

- 13.** A population has 30% who are smokers with a constant force of mortality 0.2 and 70% who are non-smokers with a constant force of mortality 0.1.

Calculate the 75th percentile of the distribution of the future lifetime of an individual selected at random from this population.

- (A) 10.7
- (B) 11.0
- (C) 11.2
- (D) 11.6
- (E) 11.8

- 14.** Aggregate losses for a portfolio of policies are modeled as follows:

- (i) The number of losses before any coverage modifications follows a Poisson distribution with mean λ .
- (ii) The severity of each loss before any coverage modifications is uniformly distributed between 0 and b .

The insurer would like to model the impact of imposing an ordinary deductible, d ($0 < d < b$), on each loss and reimbursing only a percentage, c ($0 < c \leq 1$), of each loss in excess of the deductible.

It is assumed that the coverage modifications will not affect the loss distribution. The insurer models its claims with modified frequency and severity distributions. The modified claim amount is uniformly distributed on the interval $[0, c(b-d)]$.

Determine the mean of the modified frequency distribution.

- (A) λ
- (B) λc
- (C) $\lambda \frac{d}{b}$
- (D) $\lambda \frac{b-d}{b}$
- (E) $\lambda c \frac{b-d}{b}$

15. The RIP Life Insurance Company specializes in selling a fully discrete whole life insurance of 10,000 to 65 year olds by telephone. For each policy:

- (i) The annual contract premium is 500.
- (ii) Mortality follows the Illustrative Life Table.
- (iii) $i = 0.06$

The number of telephone inquiries RIP receives follows a Poisson process with mean 50 per day. 20% of the inquiries result in the sale of a policy.

The number of inquiries and the future lifetimes of all the insureds who purchase policies on a particular day are independent.

Using the normal approximation, calculate the probability that S , the total prospective loss at issue for all the policies sold on a particular day, will be less than zero.

- (A) 0.33
- (B) 0.50
- (C) 0.67
- (D) 0.84
- (E) 0.99

16. For a special fully discrete whole life insurance on (40):

- (i) The death benefit is 1000 for the first 20 years; 5000 for the next 5 years; 1000 thereafter.
- (ii) The annual benefit premium is $1000P_{40}$ for the first 20 years; $5000P_{40}$ for the next 5 years; π thereafter.
- (iii) Mortality follows the Illustrative Life Table.
- (iv) $i = 0.06$

Calculate ${}_{21}V$, the benefit reserve at the end of year 21 for this insurance.

- (A) 255
- (B) 259
- (C) 263
- (D) 267
- (E) 271

17. For a whole life insurance of 1 on (41) with death benefit payable at the end of year of death, you are given:

(i) $i = 0.05$

(ii) $p_{40} = 0.9972$

(iii) $A_{41} - A_{40} = 0.00822$

(iv) ${}^2A_{41} - {}^2A_{40} = 0.00433$

(v) Z is the present-value random variable for this insurance.

Calculate $\text{Var}(Z)$.

(A) 0.023

(B) 0.024

(C) 0.025

(D) 0.026

(E) 0.027

18. For a perpetuity-immediate with annual payments of 1:

- (i) The sequence of annual discount factors follows a Markov chain with the following three states:

State number	0	1	2
Annual discount factor, v	0.95	0.94	0.93

- (ii) The transition matrix for the annual discount factors is:

$$\begin{bmatrix} 0.0 & 1.0 & 0.0 \\ 0.9 & 0.0 & 0.1 \\ 0.0 & 1.0 & 0.0 \end{bmatrix}$$

Y is the present value of the perpetuity payments when the initial state is 1.

Calculate $E(Y)$.

- (A) 15.67
(B) 15.71
(C) 15.75
(D) 16.82
(E) 16.86

- 19.** A member of a high school math team is practicing for a contest. Her advisor has given her three practice problems: #1, #2, and #3.

She randomly chooses one of the problems, and works on it until she solves it. Then she randomly chooses one of the remaining unsolved problems, and works on it until solved. Then she works on the last unsolved problem.

She solves problems at a Poisson rate of 1 problem per 5 minutes.

Calculate the probability that she has solved problem #3 within 10 minutes of starting the problems.

- (A) 0.18
- (B) 0.34
- (C) 0.45
- (D) 0.51
- (E) 0.59

- 20.** For a double decrement table, you are given:

(i) $\mu_x^{(1)}(t) = 0.2 \mu_x^{(\tau)}(t), \quad t > 0$

(ii) $\mu_x^{(\tau)}(t) = k t^2, \quad t > 0$

(iii) $q_x^{(1)} = 0.04$

Calculate ${}_2q_x^{(2)}$.

- (A) 0.45
- (B) 0.53
- (C) 0.58
- (D) 0.64
- (E) 0.73

21. For (x) :

(i) K is the curtate future lifetime random variable.

(ii) $q_{x+k} = 0.1(k+1)$, $k = 0, 1, 2, \dots, 9$

Calculate $\text{Var}(K \wedge 3)$.

(A) 1.1


(B) 1.2

(C) 1.3

(D) 1.4

(E) 1.5

22. The graph of the density function for losses is:



Calculate the loss elimination ratio for an ordinary deductible of 20.

(A) 0.20

(B) 0.24

(C) 0.28

(D) 0.32

(E) 0.36

23. Michel, age 45, is expected to experience higher than standard mortality only at age 64. For a special fully discrete whole life insurance of 1 on Michel, you are given:

- (i) The benefit premiums are not level.
- (ii) The benefit premium for year 20, π_{19} , exceeds P_{45} for a standard risk by 0.010.
- (iii) Benefit reserves on his insurance are the same as benefit reserves for a fully discrete whole life insurance of 1 on (45) with standard mortality and level benefit premiums.
- (iv) $i = 0.03$
- (v) ${}_{20}V_{45} = 0.427$

Calculate the excess of q_{64} for Michel over the standard q_{64} .

- (A) 0.012
- (B) 0.014
- (C) 0.016
- (D) 0.018
- (E) 0.020

24. For a block of fully discrete whole life insurances of 1 on independent lives age x , you are given:

(i) $i = 0.06$

(ii) $A_x = 0.24905$

(iii) ${}^2A_x = 0.09476$

(iv) $\pi = 0.025$, where π is the contract premium for each policy.

(v) Losses are based on the contract premium.

Using the normal approximation, calculate the minimum number of policies the insurer must issue so that the probability of a positive total loss on the policies issued is less than or equal to 0.05.

(A) 25

(B) 27

(C) 29

(D) 31

(E) 33

- 25.** Your company currently offers a whole life annuity product that pays the annuitant 12,000 at the beginning of each year. A member of your product development team suggests enhancing the product by adding a death benefit that will be paid at the end of the year of death.

Using a discount rate, d , of 8%, calculate the death benefit that minimizes the variance of the present value random variable of the new product.

- (A) 0
- (B) 50,000
- (C) 100,000
- (D) 150,000
- (E) 200,000

- 26.** A towing company provides all towing services to members of the City Automobile Club. You are given:

Towing Distance	Towing Cost	Frequency
0-9.99 miles	80	50%
10-29.99 miles	100	40%
30+ miles	160	10%

- (i) The automobile owner must pay 10% of the cost and the remainder is paid by the City Automobile Club.
- (ii) The number of towings has a Poisson distribution with mean of 1000 per year.
- (iii) The number of towings and the costs of individual towings are all mutually independent.

Using the normal approximation for the distribution of aggregate towing costs, calculate the probability that the City Automobile Club pays more than 90,000 in any given year.

- (A) 3%
- (B) 10%
- (C) 50%
- (D) 90%
- (E) 97%

27. You are given:

- (i) Losses follow an exponential distribution with the same mean in all years.
- (ii) The loss elimination ratio this year is 70%.
- (iii) The ordinary deductible for the coming year is $4/3$ of the current deductible.

Compute the loss elimination ratio for the coming year.

- (A) 70%
- (B) 75%
- (C) 80%
- (D) 85%
- (E) 90%

28. For T , the future lifetime random variable for (0):

- (i) $\omega > 70$
- (ii) ${}_{40}P_0 = 0.6$
- (iii) $E(T) = 62$
- (iv) $E(T \wedge t) = t - 0.005t^2, \quad 0 < t < 60$

Calculate the complete expectation of life at 40.

- (A) 30
- (B) 35
- (C) 40
- (D) 45
- (E) 50

- 29.** Two actuaries use the same mortality table to price a fully discrete 2-year endowment insurance of 1000 on (x) .
- (i) Kevin calculates non-level benefit premiums of 608 for the first year and 350 for the second year.
 - (ii) Kira calculates level annual benefit premiums of π .
 - (iii) $d = 0.05$

Calculate π .

- (A) 482
- (B) 489
- (C) 497
- (D) 508
- (E) 517

30. For a fully discrete 10-payment whole life insurance of 100,000 on (x) , you are given:

- (i) $i = 0.05$
- (ii) $q_{x+9} = 0.011$
- (iii) $q_{x+10} = 0.012$
- (iv) $q_{x+11} = 0.014$
- (v) The level annual benefit premium is 2078.
- (vi) The benefit reserve at the end of year 9 is 32,535.

Calculate $100,000A_{x+11}$.

- (A) 34,100
- (B) 34,300
- (C) 35,500
- (D) 36,500
- (E) 36,700

31. You are given:

- (i) Mortality follows DeMoivre's law with $\omega = 105$.
- (ii) (45) and (65) have independent future lifetimes.

Calculate $\overset{\circ}{e}_{45:65}$.

- (A) 33
- (B) 34
- (C) 35
- (D) 36
- (E) 37

32. Given: The survival function $s(x)$, where

$$s(x) = 1, \quad 0 \leq x < 1$$

$$s(x) = 1 - \left\{ \frac{e^x}{100} \right\}, \quad 1 \leq x < 4.5$$

$$s(x) = 0, \quad 4.5 \leq x$$

Calculate $\mu(4)$.

- (A) 0.45
- (B) 0.55
- (C) 0.80
- (D) 1.00
- (E) 1.20

33. For a triple decrement table, you are given:

(i) $\mu_x^{(1)}(t) = 0.3, \quad t > 0$

(ii) $\mu_x^{(2)}(t) = 0.5, \quad t > 0$

(iii) $\mu_x^{(3)}(t) = 0.7, \quad t > 0$

Calculate $q_x^{(2)}$.

- (A) 0.26
- (B) 0.30
- (C) 0.33
- (D) 0.36
- (E) 0.39

34. You are given:

(i) the following select-and-ultimate mortality table with 3-year select period:

x	$q_{[x]}$	$q_{[x]+1}$	$q_{[x]+2}$	q_{x+3}	$x+3$
60	0.09	0.11	0.13	0.15	63
61	0.10	0.12	0.14	0.16	64
62	0.11	0.13	0.15	0.17	65
63	0.12	0.14	0.16	0.18	66
64	0.13	0.15	0.17	0.19	67

(ii) $i = 0.03$

Calculate ${}_{2|2}A_{[60]}$, the actuarial present value of a 2-year deferred 2-year term insurance on $[60]$.

- (A) 0.156
- (B) 0.160
- (C) 0.186
- (D) 0.190
- (E) 0.195

35. You are given:

(i) $\mu_x(t) = 0.01, \quad 0 \leq t < 5$

(ii) $\mu_x(t) = 0.02, \quad 5 \leq t$

(iii) $\delta = 0.06$

Calculate \bar{a}_x .

(A) 12.5

(B) 13.0

(C) 13.4

(D) 13.9

(E) 14.3

36. Actuaries have modeled auto windshield claim frequencies. They have concluded that the number of windshield claims filed per year per driver follows the Poisson distribution with parameter λ , where λ follows the gamma distribution with mean 3 and variance 3.

Calculate the probability that a driver selected at random will file no more than 1 windshield claim next year.

(A) 0.15

(B) 0.19

(C) 0.20

(D) 0.24

(E) 0.31

- 37.** The number of auto vandalism claims reported per month at Sunny Daze Insurance Company (SDIC) has mean 110 and variance 750. Individual losses have mean 1101 and standard deviation 70. The number of claims and the amounts of individual losses are independent.

Using the normal approximation, calculate the probability that SDIC's aggregate auto vandalism losses reported for a month will be less than 100,000.

- (A) 0.24
- (B) 0.31
- (C) 0.36
- (D) 0.39
- (E) 0.49

- 38.** For an allosaur with 10,000 calories stored at the start of a day:

- (i) The allosaur uses calories uniformly at a rate of 5,000 per day. If his stored calories reach 0, he dies.
- (ii) Each day, the allosaur eats 1 scientist (10,000 calories) with probability 0.45 and no scientist with probability 0.55.
- (iii) The allosaur eats only scientists.
- (iv) The allosaur can store calories without limit until needed.

Calculate the probability that the allosaur ever has 15,000 or more calories stored.

- (A) 0.54
- (B) 0.57
- (C) 0.60
- (D) 0.63
- (E) 0.66

39. Lucky Tom finds coins on his way to work at a Poisson rate of 0.5 coins/minute. The denominations are randomly distributed:

- (i) 60% of the coins are worth 1 each
- (ii) 20% of the coins are worth 5 each
- (iii) 20% of the coins are worth 10 each.

Calculate the probability that in the first ten minutes of his walk he finds at least 2 coins worth 10 each, and in the first twenty minutes finds at least 3 coins worth 10 each.

- (A) 0.08
- (B) 0.12
- (C) 0.16
- (D) 0.20
- (E) 0.24

40. For a fully discrete whole life insurance of 1000 on (60), the annual benefit premium was calculated using the following:

(i) $i = 0.06$

(ii) $q_{60} = 0.01376$

(iii) $1000A_{60} = 369.33$

(iv) $1000A_{61} = 383.00$

A particular insured is expected to experience a first-year mortality rate ten times the rate used to calculate the annual benefit premium. The expected mortality rates for all other years are the ones originally used.

Calculate the expected loss at issue for this insured, based on the original benefit premium.

(A) 72

(B) 86

(C) 100

(D) 114

(E) 128

41. For a fully discrete whole life insurance of 1000 on (40), you are given:

(i) $i = 0.06$

(ii) Mortality follows the Illustrative Life Table.

(iii) $\ddot{a}_{40:\overline{10}|} = 7.70$

(iv) $\ddot{a}_{50:\overline{10}|} = 7.57$

(v) $1000A^1_{40:\overline{20}|} = 60.00$

At the end of the tenth year, the insured elects an option to retain the coverage of 1000 for life, but pay premiums for the next ten years only.

Calculate the revised annual benefit premium for the next 10 years.

(A) 11

(B) 15

(C) 17

(D) 19

(E) 21

42. For a double-decrement table where cause 1 is death and cause 2 is withdrawal, you are given:

- (i) Deaths are uniformly distributed over each year of age in the single-decrement table.
- (ii) Withdrawals occur only at the end of each year of age.
- (iii) $l_x^{(\tau)} = 1000$
- (iv) $q_x^{(2)} = 0.40$
- (v) $d_x^{(1)} = 0.45 d_x^{(2)}$

Calculate $p_x^{(2)}$.

- (A) 0.51
- (B) 0.53
- (C) 0.55
- (D) 0.57
- (E) 0.59

43. You intend to hire 200 employees for a new management-training program. To predict the number who will complete the program, you build a multiple decrement table. You decide that the following associated single decrement assumptions are appropriate:

- (i) Of 40 hires, the number who fail to make adequate progress in each of the first three years is 10, 6, and 8, respectively.
- (ii) Of 30 hires, the number who resign from the company in each of the first three years is 6, 8, and 2, respectively.
- (iii) Of 20 hires, the number who leave the program for other reasons in each of the first three years is 2, 2, and 4, respectively.
- (iv) You use the uniform distribution of decrements assumption in each year in the multiple decrement table.

Calculate the expected number who fail to make adequate progress in the third year.

- (A) 4
- (B) 8
- (C) 12
- (D) 14
- (E) 17

- 44.** Bob is an overworked underwriter. Applications arrive at his desk at a Poisson rate of 60 per day. Each application has a $1/3$ chance of being a “bad” risk and a $2/3$ chance of being a “good” risk.

Since Bob is overworked, each time he gets an application he flips a fair coin. If it comes up heads, he accepts the application without looking at it. If the coin comes up tails, he accepts the application if and only if it is a “good” risk. The expected profit on a “good” risk is 300 with variance 10,000. The expected profit on a “bad” risk is -100 with variance 90,000.

Calculate the variance of the profit on the applications he accepts today.

- (A) 4,000,000
- (B) 4,500,000
- (C) 5,000,000
- (D) 5,500,000
- (E) 6,000,000

- 45.** Prescription drug losses, S , are modeled assuming the number of claims has a geometric distribution with mean 4, and the amount of each prescription is 40.

Calculate $E[(S - 100)_+]$.

- (A) 60
- (B) 82
- (C) 92
- (D) 114
- (E) 146

46. For a temporary life annuity-immediate on independent lives (30) and (40):

(i) Mortality follows the Illustrative Life Table.

(ii) $i = 0.06$

Calculate $a_{30:40:\overline{10}|}$.

(A) 6.64

(B) 7.17

(C) 7.88

(D) 8.74

(E) 9.86

47. For a special whole life insurance on (35), you are given:

(i) The annual benefit premium is payable at the beginning of each year.

(ii) The death benefit is equal to 1000 plus the return of all benefit premiums paid in the past without interest.

(iii) The death benefit is paid at the end of the year of death.

(iv) $A_{35} = 0.42898$

(v) $(IA)_{35} = 6.16761$

(vi) $i = 0.05$

Calculate the annual benefit premium for this insurance.

(A) 73.66

(B) 75.28

(C) 77.42

(D) 78.95

(E) 81.66

- 48.** Subway trains arrive at a station at a Poisson rate of 20 per hour. 25% of the trains are express and 75% are local. The types of each train are independent. An express gets you to work in 16 minutes and a local gets you there in 28 minutes. You always take the first train to arrive. Your co-worker always takes the first express. You both are waiting at the same station.

Which of the following is true?

- (A) Your expected arrival time is 6 minutes earlier than your co-worker's.
- (B) Your expected arrival time is 4.5 minutes earlier than your co-worker's.
- (C) Your expected arrival times are the same.
- (D) Your expected arrival time is 4.5 minutes later than your co-worker's.
- (E) Your expected arrival time is 6 minutes later than your co-worker's.

- 49.** For a special fully continuous whole life insurance of 1 on the last-survivor of (x) and (y) , you are given:

- (i) $T(x)$ and $T(y)$ are independent.
- (ii) $\mu_x(t) = \mu_y(t) = 0.07, \quad t > 0$
- (iii) $\delta = 0.05$
- (iv) Premiums are payable until the first death.

Calculate the level annual benefit premium for this insurance.

- (A) 0.04
- (B) 0.07
- (C) 0.08
- (D) 0.10
- (E) 0.14

50. For a fully discrete whole life insurance of 1000 on (20), you are given:

- (i) $1000 P_{20} = 10$
- (ii) $1000 {}_{20}V_{20} = 490$
- (iii) $1000 {}_{21}V_{20} = 545$
- (iv) $1000 {}_{22}V_{20} = 605$
- (v) $q_{40} = 0.022$

Calculate q_{41} .

- (A) 0.024
- (B) 0.025
- (C) 0.026
- (D) 0.027
- (E) 0.028

51. For a fully discrete whole life insurance of 1000 on (60), you are given:

- (i) $i = 0.06$
- (ii) Mortality follows the Illustrative Life Table, except that there are extra mortality risks at age 60 such that $q_{60} = 0.015$.

Calculate the annual benefit premium for this insurance.

- (A) 31.5
- (B) 32.0
- (C) 32.1
- (D) 33.1
- (E) 33.2

- 52.** At the beginning of each round of a game of chance the player pays 12.5. The player then rolls one die with outcome N . The player then rolls N dice and wins an amount equal to the total of the numbers showing on the N dice. All dice have 6 sides and are fair.

Using the normal approximation, calculate the probability that a player starting with 15,000 will have at least 15,000 after 1000 rounds.

- (A) 0.01
- (B) 0.04
- (C) 0.06
- (D) 0.09
- (E) 0.12

- 53.** X is a discrete random variable with a probability function which is a member of the $(a,b,0)$ class of distributions.

You are given:

- (i) $P(X = 0) = P(X = 1) = 0.25$
- (ii) $P(X = 2) = 0.1875$

Calculate $P(X = 3)$.

- (A) 0.120
- (B) 0.125
- (C) 0.130
- (D) 0.135
- (E) 0.140

54. Nancy reviews the interest rates each year for a 30-year fixed mortgage issued on July 1. She models interest rate behavior by a Markov model assuming:

- (i) Interest rates always change between years.
- (ii) The change in any given year is dependent on the change in prior years as follows:

from year $t - 3$ to year $t - 2$	from year $t - 2$ to year $t - 1$	Probability that year t will increase from year $t - 1$
Increase	Increase	0.10
Decrease	Decrease	0.20
Increase	Decrease	0.40
Decrease	Increase	0.25

She notes that interest rates decreased from year 2000 to 2001 and from year 2001 to 2002.

Calculate the probability that interest rates will decrease from year 2003 to 2004.

- (A) 0.76
- (B) 0.79
- (C) 0.82
- (D) 0.84
- (E) 0.87

55. For a 20-year deferred whole life annuity-due of 1 per year on (45), you are given:

(i) Mortality follows De Moivre's law with $\omega = 105$.

(ii) $i = 0$

Calculate the probability that the sum of the annuity payments actually made will exceed the actuarial present value at issue of the annuity.

(A) 0.425

(B) 0.450

(C) 0.475

(D) 0.500

(E) 0.525

56. For a continuously increasing whole life insurance on (x) , you are given:

(i) The force of mortality is constant.

(ii) $\delta = 0.06$

(iii) ${}^2\bar{A}_x = 0.25$

Calculate $(\bar{IA})_x$.

(A) 2.889

(B) 3.125

(C) 4.000

(D) 4.667

(E) 5.500

- 57.** XYZ Co. has just purchased two new tools with independent future lifetimes. Each tool has its own distinct De Moivre survival pattern. One tool has a 10-year maximum lifetime and the other a 7-year maximum lifetime.

Calculate the expected time until both tools have failed.

- (A) 5.0
- (B) 5.2
- (C) 5.4
- (D) 5.6
- (E) 5.8

- 58.** XYZ Paper Mill purchases a 5-year special insurance paying a benefit in the event its machine breaks down. If the cause is “minor” (1), only a repair is needed. If the cause is “major” (2), the machine must be replaced.

Given:

- (i) The benefit for cause (1) is 2000 payable at the moment of breakdown.
- (ii) The benefit for cause (2) is 500,000 payable at the moment of breakdown.
- (iii) Once a benefit is paid, the insurance contract is terminated.
- (iv) $\mu^{(1)}(t) = 0.100$ and $\mu^{(2)}(t) = 0.004$, for $t > 0$
- (v) $\delta = 0.04$

Calculate the actuarial present value of this insurance.

- (A) 7840
- (B) 7880
- (C) 7920
- (D) 7960
- (E) 8000

59. You are given:

(i) $R = 1 - e^{-\int_0^1 \mu_x(t) dt}$

(ii) $S = 1 - e^{-\int_0^1 (\mu_x(t) + k) dt}$

(iii) k is a constant such that $S = 0.75R$

Determine an expression for k .

(A) $\ln((1 - q_x) / (1 - 0.75q_x))$

(B) $\ln((1 - 0.75q_x) / (1 - p_x))$

(C) $\ln((1 - 0.75p_x) / (1 - p_x))$

(D) $\ln((1 - p_x) / (1 - 0.75q_x))$

(E) $\ln((1 - 0.75q_x) / (1 - q_x))$

60. The number of claims in a period has a geometric distribution with mean 4. The amount of each claim X follows $P(X = x) = 0.25$, $x = 1, 2, 3, 4$. The number of claims and the claim amounts are independent. S is the aggregate claim amount in the period.

Calculate $F_s(3)$.

(A) 0.27

(B) 0.29

(C) 0.31

(D) 0.33

(E) 0.35

- 61.** Insurance agent Hunt N. Quotum will receive no annual bonus if the ratio of incurred losses to earned premiums for his book of business is 60% or more for the year. If the ratio is less than 60%, Hunt's bonus will be a percentage of his earned premium equal to 15% of the difference between his ratio and 60%. Hunt's annual earned premium is 800,000.

Incurred losses are distributed according to the Pareto distribution, with $\theta = 500,000$ and $\alpha = 2$.

Calculate the expected value of Hunt's bonus.

- (A) 13,000
- (B) 17,000
- (C) 24,000
- (D) 29,000
- (E) 35,000

- 62.** A large machine in the ABC Paper Mill is 25 years old when ABC purchases a 5-year term insurance paying a benefit in the event the machine breaks down.

Given:

- (i) Annual benefit premiums of 6643 are payable at the beginning of the year.
- (ii) A benefit of 500,000 is payable at the moment of breakdown.
- (iii) Once a benefit is paid, the insurance contract is terminated.
- (iv) Machine breakdowns follow De Moivre's law with $l_x = 100 - x$.
- (v) $i = 0.06$

Calculate the benefit reserve for this insurance at the end of the third year.

- (A) -91
- (B) 0
- (C) 163
- (D) 287
- (E) 422

63. For a whole life insurance of 1 on (x) , you are given:

- (i) The force of mortality is $\mu_x(t)$.
- (ii) The benefits are payable at the moment of death.
- (iii) $\delta = 0.06$
- (iv) $\bar{A}_x = 0.60$

Calculate the revised actuarial present value of this insurance assuming $\mu_x(t)$ is increased by 0.03 for all t and δ is decreased by 0.03.

- (A) 0.5
- (B) 0.6
- (C) 0.7
- (D) 0.8
- (E) 0.9

64. A maintenance contract on a hotel promises to replace burned out light bulbs at the end of each year for three years. The hotel has 10,000 light bulbs. The light bulbs are all new. If a replacement bulb burns out, it too will be replaced with a new bulb.

You are given:

- (i) For new light bulbs, $q_0 = 0.10$
 $q_1 = 0.30$
 $q_2 = 0.50$

(ii) Each light bulb costs 1.

(iii) $i = 0.05$

Calculate the actuarial present value of this contract.

- (A) 6700
- (B) 7000
- (C) 7300
- (D) 7600
- (E) 8000

65. You are given:

$$\mu(x) = \begin{cases} 0.04, & 0 < x < 40 \\ 0.05, & x > 40 \end{cases}$$

Calculate $\overset{\circ}{e}_{25:\overline{25}|}$.

- (A) 14.0
- (B) 14.4
- (C) 14.8
- (D) 15.2
- (E) 15.6

66. For a select-and-ultimate mortality table with a 3-year select period:

(i)

x	$q_{[x]}$	$q_{[x]+1}$	$q_{[x]+2}$	q_{x+3}	$x+3$
60	0.09	0.11	0.13	0.15	63
61	0.10	0.12	0.14	0.16	64
62	0.11	0.13	0.15	0.17	65
63	0.12	0.14	0.16	0.18	66
64	0.13	0.15	0.17	0.19	67

- (ii) White was a newly selected life on 01/01/2000.
- (iii) White's age on 01/01/2001 is 61.
- (iv) P is the probability on 01/01/2001 that White will be alive on 01/01/2006.

Calculate P .

- (A) $0 \leq P < 0.43$
- (B) $0.43 \leq P < 0.45$
- (C) $0.45 \leq P < 0.47$
- (D) $0.47 \leq P < 0.49$
- (E) $0.49 \leq P \leq 1.00$

- 67.** For a continuous whole life annuity of 1 on (x) :
- (i) $T(x)$ is the future lifetime random variable for (x) .
 - (ii) The force of interest and force of mortality are constant and equal.
 - (iii) $\bar{a}_x = 12.50$

Calculate the standard deviation of $\bar{a}_{\overline{T(x)|}}$.

- (A) 1.67
- (B) 2.50
- (C) 2.89
- (D) 6.25
- (E) 7.22

- 68.** For a special fully discrete whole life insurance on (x) :

- (i) The death benefit is 0 in the first year and 5000 thereafter.
- (ii) Level benefit premiums are payable for life.
- (iii) $q_x = 0.05$
- (iv) $v = 0.90$
- (v) $\ddot{a}_x = 5.00$
- (vi) ${}_{10}V_x = 0.20$
- (vii) ${}_{10}V$ is the benefit reserve at the end of year 10 for this insurance.

Calculate ${}_{10}V$.

- (A) 795
- (B) 1000
- (C) 1090
- (D) 1180
- (E) 1225

- 69.** For a fully discrete 2-year term insurance of 1 on (x) :
- (i) 0.95 is the lowest premium such that there is a 0% chance of loss in year 1.
 - (ii) $p_x = 0.75$
 - (iii) $p_{x+1} = 0.80$
 - (iv) Z is the random variable for the present value at issue of future benefits.

Calculate $\text{Var}(Z)$.

- (A) 0.15
 - (B) 0.17
 - (C) 0.19
 - (D) 0.21
 - (E) 0.23
- 70.** A group dental policy has a negative binomial claim count distribution with mean 300 and variance 800.

Ground-up severity is given by the following table:

Severity	Probability
40	0.25
80	0.25
120	0.25
200	0.25

You expect severity to increase 50% with no change in frequency. You decide to impose a per claim deductible of 100.

Calculate the expected total claim payment after these changes.

- (A) Less than 18,000
- (B) At least 18,000, but less than 20,000
- (C) At least 20,000, but less than 22,000
- (D) At least 22,000, but less than 24,000
- (E) At least 24,000

- 71.** You own a fancy light bulb factory. Your workforce is a bit clumsy – they keep dropping boxes of light bulbs. The boxes have varying numbers of light bulbs in them, and when dropped, the entire box is destroyed.

You are given:

Expected number of boxes dropped per month:	50
Variance of the number of boxes dropped per month:	100
Expected value per box:	200
Variance of the value per box:	400

You pay your employees a bonus if the value of light bulbs destroyed in a month is less than 8000.

Assuming independence and using the normal approximation, calculate the probability that you will pay your employees a bonus next month.

- (A) 0.16
- (B) 0.19
- (C) 0.23
- (D) 0.27
- (E) 0.31

- 72.** Each of 100 independent lives purchase a single premium 5-year deferred whole life insurance of 10 payable at the moment of death. You are given:

- (i) $\mu = 0.04$
- (ii) $\delta = 0.06$
- (iii) F is the aggregate amount the insurer receives from the 100 lives.

Using the normal approximation, calculate F such that the probability the insurer has sufficient funds to pay all claims is 0.95.

- (A) 280
- (B) 390
- (C) 500
- (D) 610
- (E) 720

73. For a select-and-ultimate table with a 2-year select period:

x	$P_{[x]}$	$P_{[x]+1}$	P_{x+2}	$x+2$
48	0.9865	0.9841	0.9713	50
49	0.9858	0.9831	0.9698	51
50	0.9849	0.9819	0.9682	52
51	0.9838	0.9803	0.9664	53

Keith and Clive are independent lives, both age 50. Keith was selected at age 45 and Clive was selected at age 50.

Calculate the probability that exactly one will be alive at the end of three years.

- (A) Less than 0.115
- (B) At least 0.115, but less than 0.125
- (C) At least 0.125, but less than 0.135
- (D) At least 0.135, but less than 0.145
- (E) At least 0.145

74-75. Use the following information for questions 74 and 75.

For a tyrannosaur with 10,000 calories stored:

- (i) The tyrannosaur uses calories uniformly at a rate of 10,000 per day. If his stored calories reach 0, he dies.
- (ii) The tyrannosaur eats scientists (10,000 calories each) at a Poisson rate of 1 per day.
- (iii) The tyrannosaur eats only scientists.
- (iv) The tyrannosaur can store calories without limit until needed.

74. Calculate the probability that the tyrannosaur dies within the next 2.5 days.

- (A) 0.30
- (B) 0.40
- (C) 0.50
- (D) 0.60
- (E) 0.70

75. Calculate the expected calories eaten in the next 2.5 days.

- (A) 17,800
- (B) 18,800
- (C) 19,800
- (D) 20,800
- (E) 21,800

76. A fund is established by collecting an amount P from each of 100 independent lives age 70. The fund will pay the following benefits:

- 10, payable at the end of the year of death, for those who die before age 72, or
- P , payable at age 72, to those who survive.

You are given:

Mortality follows the Illustrative Life Table.

(i) $i = 0.08$

Calculate P , using the equivalence principle.

- (A) 2.33
- (B) 2.38
- (C) 3.02
- (D) 3.07
- (E) 3.55

77. You are given:

(i) $P_x = 0.090$

(ii) ${}_nV_x = 0.563$

(iii) $P_{x:\overline{n}|}^1 = 0.00864$

Calculate $P_{x:\overline{n}|}^1$.

- (A) 0.008
- (B) 0.024
- (C) 0.040
- (D) 0.065
- (E) 0.085

78. You are given:

(i) Mortality follows De Moivre's law with $\omega = 100$.

(ii) $i = 0.05$

(iii) The following annuity-certain values:

$$\bar{a}_{40|} = 17.58$$

$$\bar{a}_{50|} = 18.71$$

$$\bar{a}_{60|} = 19.40$$

Calculate ${}_{10}\bar{V}(\bar{A}_{40})$.

(A) 0.075

(B) 0.077

(C) 0.079

(D) 0.081

(E) 0.083

79. For a group of individuals all age x , you are given:

(i) 30% are smokers and 70% are non-smokers.

(ii) The constant force of mortality for smokers is 0.06.

(iii) The constant force of mortality for non-smokers is 0.03.

(iv) $\delta = 0.08$

Calculate $\text{Var}\left(\bar{a}_{T(x)|}\right)$ for an individual chosen at random from this group.

(A) 13.0

(B) 13.3

(C) 13.8

(D) 14.1

(E) 14.6

- 80.** For a certain company, losses follow a Poisson frequency distribution with mean 2 per year, and the amount of a loss is 1, 2, or 3, each with probability $1/3$. Loss amounts are independent of the number of losses, and of each other.

An insurance policy covers all losses in a year, subject to an annual aggregate deductible of 2.

Calculate the expected claim payments for this insurance policy.

- (A) 2.00
- (B) 2.36
- (C) 2.45
- (D) 2.81
- (E) 2.96

- 81.** A Poisson claims process has two types of claims, Type I and Type II.

- (i) The expected number of claims is 3000.
- (ii) The probability that a claim is Type I is $1/3$.
- (iii) Type I claim amounts are exactly 10 each.
- (iv) The variance of aggregate claims is 2,100,000.

Calculate the variance of aggregate claims with Type I claims excluded.

- (A) 1,700,000
- (B) 1,800,000
- (C) 1,900,000
- (D) 2,000,000
- (E) 2,100,000

82. Don, age 50, is an actuarial science professor. His career is subject to two decrements:

- (i) Decrement 1 is mortality. The associated single decrement table follows De Moivre's law with $\omega = 100$.
- (ii) Decrement 2 is leaving academic employment, with

$$\mu_{50}^{(2)}(t) = 0.05, \quad t \geq 0$$

Calculate the probability that Don remains an actuarial science professor for at least five but less than ten years.

- (A) 0.22
- (B) 0.25
- (C) 0.28
- (D) 0.31
- (E) 0.34

83. For a double decrement model:

- (i) In the single decrement table associated with cause (1), $q_{40}^{(1)} = 0.100$ and decrements are uniformly distributed over the year.
- (ii) In the single decrement table associated with cause (2), $q_{40}^{(2)} = 0.125$ and all decrements occur at time 0.7.

Calculate $q_{40}^{(2)}$.

- (A) 0.114
- (B) 0.115
- (C) 0.116
- (D) 0.117
- (E) 0.118

- 84.** For a special 2-payment whole life insurance on (80):
- (i) Premiums of π are paid at the beginning of years 1 and 3.
 - (ii) The death benefit is paid at the end of the year of death.
 - (iii) There is a partial refund of premium feature:
 - If (80) dies in either year 1 or year 3, the death benefit is $1000 + \frac{\pi}{2}$.
 - Otherwise, the death benefit is 1000.
 - (iv) Mortality follows the Illustrative Life Table.
 - (v) $i = 0.06$

Calculate π , using the equivalence principle.

- (A) 369
- (B) 381
- (C) 397
- (D) 409
- (E) 425

- 85.** For a special fully continuous whole life insurance on (65):

- (i) The death benefit at time t is $b_t = 1000e^{0.04t}$, $t \geq 0$.
- (ii) Level benefit premiums are payable for life.
- (iii) $\mu_{65}(t) = 0.02$, $t \geq 0$
- (iv) $\delta = 0.04$

Calculate ${}_2\bar{V}$, the benefit reserve at the end of year 2.

- (A) 0
- (B) 29
- (C) 37
- (D) 61
- (E) 83

86. You are given:

(i) $A_x = 0.28$

(ii) $A_{x+20} = 0.40$

(iii) $A_{\overline{x:20}|} = 0.25$

(iv) $i = 0.05$

Calculate $a_{\overline{x:20}|}$.

(A) 11.0

(B) 11.2

(C) 11.7

(D) 12.0

(E) 12.3

87. On his walk to work, Lucky Tom finds coins on the ground at a Poisson rate. The Poisson rate, expressed in coins per minute, is constant during any one day, but varies from day to day according to a gamma distribution with mean 2 and variance 4.

Calculate the probability that Lucky Tom finds exactly one coin during the sixth minute of today's walk.

(A) 0.22

(B) 0.24

(C) 0.26

(D) 0.28

(E) 0.30

- 88.** The unlimited severity distribution for claim amounts under an auto liability insurance policy is given by the cumulative distribution:

$$F(x) = 1 - 0.8e^{-0.02x} - 0.2e^{-0.001x}, \quad x \geq 0$$

The insurance policy pays amounts up to a limit of 1000 per claim.

Calculate the expected payment under this policy for one claim.

- (A) 57
- (B) 108
- (C) 166
- (D) 205
- (E) 240

- 89.** A machine is in one of four states (F, G, H, I) and migrates annually among them according to a Markov process with transition matrix:

	F	G	H	I
F	0.20	0.80	0.00	0.00
G	0.50	0.00	0.50	0.00
H	0.75	0.00	0.00	0.25
I	1.00	0.00	0.00	0.00

At time 0, the machine is in State F. A salvage company will pay 500 at the end of 3 years if the machine is in State F.

Assuming $v = 0.90$, calculate the actuarial present value at time 0 of this payment.

- (A) 150
- (B) 155
- (C) 160
- (D) 165
- (E) 170

- 90.** The claims department of an insurance company receives envelopes with claims for insurance coverage at a Poisson rate of $\lambda = 50$ envelopes per week. For any period of time, the number of envelopes and the numbers of claims in the envelopes are independent. The numbers of claims in the envelopes have the following distribution:

<u>Number of Claims</u>	<u>Probability</u>
1	0.20
2	0.25
3	0.40
4	0.15

Using the normal approximation, calculate the 90th percentile of the number of claims received in 13 weeks.

- (A) 1690
(B) 1710
(C) 1730
(D) 1750
(E) 1770

- 91.** You are given:

- (i) The survival function for males is $s(x) = 1 - \frac{x}{75}$, $0 < x < 75$.
(ii) Female mortality follows De Moivre's law.
(iii) At age 60, the female force of mortality is 60% of the male force of mortality.

For two independent lives, a male age 65 and a female age 60, calculate the expected time until the second death.

- (A) 4.33
(B) 5.63
(C) 7.23
(D) 11.88
(E) 13.17

92. For a fully continuous whole life insurance of 1:

(i) $\mu = 0.04$

(ii) $\delta = 0.08$

(iii) L is the loss-at-issue random variable based on the benefit premium.

Calculate $\text{Var}(L)$.

(A) $\frac{1}{10}$

(B) $\frac{1}{5}$

(C) $\frac{1}{4}$

(D) $\frac{1}{3}$

(E) $\frac{1}{2}$

93. The random variable for a loss, X , has the following characteristics:

x	$F(x)$	$E(X \wedge x)$
0	0.0	0
100	0.2	91
200	0.6	153
1000	1.0	331

Calculate the mean excess loss for a deductible of 100.

(A) 250

(B) 300

(C) 350

(D) 400

(E) 450

94. WidgetsRUs owns two factories. It buys insurance to protect itself against major repair costs. Profit equals revenues, less the sum of insurance premiums, retained major repair costs, and all other expenses. WidgetsRUs will pay a dividend equal to the profit, if it is positive.

You are given:

- (i) Combined revenue for the two factories is 3.
- (ii) Major repair costs at the factories are independent.
- (iii) The distribution of major repair costs for each factory is

<u>k</u>	<u>Prob (k)</u>
0	0.4
1	0.3
2	0.2
3	0.1

- (iv) At each factory, the insurance policy pays the major repair costs in excess of that factory's ordinary deductible of 1. The insurance premium is 110% of the expected claims.
- (v) All other expenses are 15% of revenues.

Calculate the expected dividend.

- (A) 0.43
- (B) 0.47
- (C) 0.51
- (D) 0.55
- (E) 0.59

- 95.** For watches produced by a certain manufacturer:
- (i) Lifetimes follow a single-parameter Pareto distribution with $\alpha > 1$ and $\theta = 4$.
 - (ii) The expected lifetime of a watch is 8 years.

Calculate the probability that the lifetime of a watch is at least 6 years.

- (A) 0.44
- (B) 0.50
- (C) 0.56
- (D) 0.61
- (E) 0.67

- 96.** For a special 3-year deferred whole life annuity-due on (x) :

- (i) $i = 0.04$
- (ii) The first annual payment is 1000.
- (iii) Payments in the following years increase by 4% per year.
- (iv) There is no death benefit during the three year deferral period.
- (v) Level benefit premiums are payable at the beginning of each of the first three years.
- (vi) $e_x = 11.05$ is the curtate expectation of life for (x) .

(vii)

k	1	2	3
${}_k P_x$	0.99	0.98	0.97

Calculate the annual benefit premium.

- (A) 2625
- (B) 2825
- (C) 3025
- (D) 3225
- (E) 3425

97. For a special fully discrete 10-payment whole life insurance on (30) with level annual benefit premium π :

(i) The death benefit is equal to 1000 plus the refund, without interest, of the benefit premiums paid.

(ii) $A_{30} = 0.102$

(iii) ${}_{10|}A_{30} = 0.088$

(iv) $(IA)_{30:\overline{10}|}^1 = 0.078$

(v) $\ddot{a}_{30:\overline{10}|} = 7.747$

Calculate π .

(A) 14.9

(B) 15.0

(C) 15.1

(D) 15.2

(E) 15.3

- 98.** For a given life age 30, it is estimated that an impact of a medical breakthrough will be an increase of 4 years in $\overset{\circ}{e}_{30}$, the complete expectation of life.

Prior to the medical breakthrough, $s(x)$ followed de Moivre's law with $\omega = 100$ as the limiting age.

Assuming de Moivre's law still applies after the medical breakthrough, calculate the new limiting age.

- (A) 104
- (B) 105
- (C) 106
- (D) 107
- (E) 108

- 99.** On January 1, 2002, Pat, age 40, purchases a 5-payment, 10-year term insurance of 100,000:

- (i) Death benefits are payable at the moment of death.
- (ii) Contract premiums of 4000 are payable annually at the beginning of each year for 5 years.
- (iii) $i = 0.05$
- (iv) L is the loss random variable at time of issue.

Calculate the value of L if Pat dies on June 30, 2004.

- (A) 77,100
- (B) 80,700
- (C) 82,700
- (D) 85,900
- (E) 88,000

100. Glen is practicing his simulation skills.

He generates 1000 values of the random variable X as follows:

- (i) He generates the observed value λ from the gamma distribution with $\alpha = 2$ and $\theta = 1$ (hence with mean 2 and variance 2).
- (ii) He then generates x from the Poisson distribution with mean λ .
- (iii) He repeats the process 999 more times: first generating a value λ , then generating x from the Poisson distribution with mean λ .
- (iv) The repetitions are mutually independent.

Calculate the expected number of times that his simulated value of X is 3.

- (A) 75
- (B) 100
- (C) 125
- (D) 150
- (E) 175

101. Lucky Tom finds coins on his way to work at a Poisson rate of 0.5 coins per minute. The denominations are randomly distributed:

- (i) 60% of the coins are worth 1;
- (ii) 20% of the coins are worth 5;
- (iii) 20% of the coins are worth 10.

Calculate the variance of the value of the coins Tom finds during his one-hour walk to work.

- (A) 379
- (B) 487
- (C) 566
- (D) 670
- (E) 768

102. For a fully discrete 20-payment whole life insurance of 1000 on (x) , you are given:

- (i) $i = 0.06$
- (ii) $q_{x+19} = 0.01254$
- (iii) The level annual benefit premium is 13.72.
- (iv) The benefit reserve at the end of year 19 is 342.03.

Calculate $1000 P_{x+20}$, the level annual benefit premium for a fully discrete whole life insurance of 1000 on $(x+20)$.

- (A) 27
- (B) 29
- (C) 31
- (D) 33
- (E) 35

103. For a multiple decrement model on (60) :

- (i) $\mu_{60}^{(1)}(t)$, $t \geq 0$, follows the Illustrative Life Table.
- (ii) $\mu_{60}^{(\tau)}(t) = 2\mu_{60}^{(1)}(t)$, $t \geq 0$

Calculate ${}_{10|}q_{60}^{(\tau)}$, the probability that decrement occurs during the 11th year.

- (A) 0.03
- (B) 0.04
- (C) 0.05
- (D) 0.06
- (E) 0.07

- 104.** (x) and (y) are two lives with identical expected mortality.
You are given:

$$P_x = P_y = 0.1$$

$P_{\overline{xy}} = 0.06$, where $P_{\overline{xy}}$ is the annual benefit premium for a fully discrete insurance of 1 on (\overline{xy}) .

$$d = 0.06$$


Calculate the premium P_{xy} , the annual benefit premium for a fully discrete insurance of 1 on (xy) .

- (A) 0.14
(B) 0.16
(C) 0.18
(D) 0.20
(E) 0.22
- 105.** For students entering a college, you are given the following from a multiple decrement model:
- (i) 1000 students enter the college at $t = 0$.
 - (ii) Students leave the college for failure (1) or all other reasons (2).
 - (iii) $\mu^{(1)}(t) = \mu \quad 0 \leq t \leq 4$
 $\mu^{(2)}(t) = 0.04 \quad 0 \leq t < 4$
 - (iv) 48 students are expected to leave the college during their first year due to all causes.

Calculate the expected number of students who will leave because of failure during their fourth year.

- (A) 8
(B) 10
(C) 24
(D) 34
(E) 41

106. The following graph is related to current human mortality:



Which of the following functions of age does the graph most likely show?

- (A) $\mu(x)$
- (B) $l_x\mu(x)$
- (C) l_xp_x
- (D) l_x
- (E) l_x^2

107. An actuary for an automobile insurance company determines that the distribution of the annual number of claims for an insured chosen at random is modeled by the negative binomial distribution with mean 0.2 and variance 0.4.

The number of claims for each individual insured has a Poisson distribution and the means of these Poisson distributions are gamma distributed over the population of insureds.

Calculate the variance of this gamma distribution.

- (A) 0.20
- (B) 0.25
- (C) 0.30
- (D) 0.35
- (E) 0.40

108. A dam is proposed for a river which is currently used for salmon breeding. You have modeled:

- (i) For each hour the dam is opened the number of salmon that will pass through and reach the breeding grounds has a distribution with mean 100 and variance 900.
- (ii) The number of eggs released by each salmon has a distribution with mean of 5 and variance of 5.
- (iii) The number of salmon going through the dam each hour it is open and the numbers of eggs released by the salmon are independent.

Using the normal approximation for the aggregate number of eggs released, determine the least number of whole hours the dam should be left open so the probability that 10,000 eggs will be released is greater than 95%.

- (A) 20
- (B) 23
- (C) 26
- (D) 29
- (E) 32

109. For a special 3-year term insurance on (x) , you are given:

- (i) Z is the present-value random variable for the death benefits.
- (ii) $q_{x+k} = 0.02(k + 1) \quad k = 0, 1, 2$
- (iii) The following death benefits, payable at the end of the year of death:

k	b_{k+1}
0	300,000
1	350,000
2	400,000

- (iv) $i = 0.06$

Calculate $E(Z)$.

- (A) 36,800
- (B) 39,100
- (C) 41,400
- (D) 43,700
- (E) 46,000

110. For a special fully discrete 20-year endowment insurance on (55):

- (i) Death benefits in year k are given by $b_k = (21 - k)$, $k = 1, 2, \dots, 20$.
- (ii) The maturity benefit is 1.
- (iii) Annual benefit premiums are level.
- (iv) ${}_kV$ denotes the benefit reserve at the end of year k , $k = 1, 2, \dots, 20$.
- (v) ${}_{10}V = 5.0$
- (vi) ${}_{19}V = 0.6$
- (vii) $q_{65} = 0.10$
- (viii) $i = 0.08$

Calculate ${}_{11}V$.

- (A) 4.5
- (B) 4.6
- (C) 4.8
- (D) 5.1
- (E) 5.3

111. For a stop-loss insurance on a three person group:

- (i) Loss amounts are independent.
- (ii) The distribution of loss amount for each person is:

<u>Loss Amount</u>	<u>Probability</u>
0	0.4
1	0.3
2	0.2
3	0.1

- (iii) The stop-loss insurance has a deductible of 1 for the group.

Calculate the net stop-loss premium.

- (A) 2.00
- (B) 2.03
- (C) 2.06
- (D) 2.09
- (E) 2.12

112. A continuous two-life annuity pays:

- 100 while both (30) and (40) are alive;
- 70 while (30) is alive but (40) is dead; and
- 50 while (40) is alive but (30) is dead.

The actuarial present value of this annuity is 1180. Continuous single life annuities paying 100 per year are available for (30) and (40) with actuarial present values of 1200 and 1000, respectively.

Calculate the actuarial present value of a two-life continuous annuity that pays 100 while at least one of them is alive.

- (A) 1400
- (B) 1500
- (C) 1600
- (D) 1700
- (E) 1800

113. For a disability insurance claim:

- (i) The claimant will receive payments at the rate of 20,000 per year, payable continuously as long as she remains disabled.
- (ii) The length of the payment period in years is a random variable with the gamma distribution with parameters $\alpha = 2$ and $\theta = 1$.
- (iii) Payments begin immediately.
- (iv) $\delta = 0.05$

Calculate the actuarial present value of the disability payments at the time of disability.

- (A) 36,400
- (B) 37,200
- (C) 38,100
- (D) 39,200
- (E) 40,000

114. For a discrete probability distribution, you are given the recursion relation

$$p(k) = \frac{2}{k} * p(k-1), \quad k = 1, 2, \dots$$

Determine $p(4)$.

- (A) 0.07
- (B) 0.08
- (C) 0.09
- (D) 0.10
- (E) 0.11

115. A company insures a fleet of vehicles. Aggregate losses have a compound Poisson distribution. The expected number of losses is 20. Loss amounts, regardless of vehicle type, have exponential distribution with $\theta = 200$.

In order to reduce the cost of the insurance, two modifications are to be made:

- (i) a certain type of vehicle will not be insured. It is estimated that this will reduce loss frequency by 20%.
- (ii) a deductible of 100 per loss will be imposed.

Calculate the expected aggregate amount paid by the insurer after the modifications.

- (A) 1600
- (B) 1940
- (C) 2520
- (D) 3200
- (E) 3880

116. For a population of individuals, you are given:

- (i) Each individual has a constant force of mortality.
- (ii) The forces of mortality are uniformly distributed over the interval $(0,2)$.

Calculate the probability that an individual drawn at random from this population dies within one year.

- (A) 0.37
- (B) 0.43
- (C) 0.50
- (D) 0.57
- (E) 0.63

- 117.** You are the producer of a television quiz show that gives cash prizes. The number of prizes, N , and prize amounts, X , have the following distributions:

n	$\Pr(N = n)$	x	$\Pr(X = x)$
1	0.8	0	0.2
2	0.2	100	0.7
		1000	0.1

Your budget for prizes equals the expected prizes plus the standard deviation of prizes.

Calculate your budget.

- (A) 306
 (B) 316
 (C) 416
 (D) 510
 (E) 518
- 118.** For a special fully discrete 3-year term insurance on (x) :
- (i) Level benefit premiums are paid at the beginning of each year.

(ii)

k	b_{k+1}	q_{x+k}
0	200,000	0.03
1	150,000	0.06
2	100,000	0.09

(iii) $i = 0.06$

Calculate the initial benefit reserve for year 2.

- (A) 6,500
 (B) 7,500
 (C) 8,100
 (D) 9,400
 (E) 10,300

119. For a special fully continuous whole life insurance on (x) :

- (i) The level premium is determined using the equivalence principle.
- (ii) Death benefits are given by $b_t = (1+i)^t$ where i is the interest rate.
- (iii) L is the loss random variable at $t = 0$ for the insurance.
- (iv) T is the future lifetime random variable of (x) .

Which of the following expressions is equal to L ?

(A) $\frac{(v^T - \bar{A}_x)}{(1 - \bar{A}_x)}$

(B) $(v^T - \bar{A}_x)(1 + \bar{A}_x)$

(C) $\frac{(v^T - \bar{A}_x)}{(1 + \bar{A}_x)}$

(D) $(v^T - \bar{A}_x)(1 - \bar{A}_x)$

(E) $\frac{(v^T + \bar{A}_x)}{(1 + \bar{A}_x)}$

120. For a 4-year college, you are given the following probabilities for dropout from all causes:

$$q_0 = 0.15$$

$$q_1 = 0.10$$

$$q_2 = 0.05$$

$$q_3 = 0.01$$

Dropouts are uniformly distributed over each year.

Compute the temporary 1.5-year complete expected college lifetime of a student entering the second year, $e_{1.5|2}^{\circ}$.

(A) 1.25

(B) 1.30

(C) 1.35

(D) 1.40

(E) 1.45

121. Lee, age 63, considers the purchase of a single premium whole life insurance of 10,000 with death benefit payable at the end of the year of death.

The company calculates benefit premiums using:

- (i) mortality based on the Illustrative Life Table,
- (ii) $i = 0.05$

The company calculates contract premiums as 112% of benefit premiums.

The single contract premium at age 63 is 5233.

Lee decides to delay the purchase for two years and invests the 5233.

Calculate the minimum annual rate of return that the investment must earn to accumulate to an amount equal to the single contract premium at age 65.

- (A) 0.030
- (B) 0.035
- (C) 0.040
- (D) 0.045
- (E) 0.050

122. You have calculated the actuarial present value of a last-survivor whole life insurance of 1 on (x) and (y) . You assumed:

- (i) The death benefit is payable at the moment of death.
- (ii) The future lifetimes of (x) and (y) are independent, and each life has a constant force of mortality with $\mu = 0.06$.
- (iii) $\delta = 0.05$

Your supervisor points out that these are not independent future lifetimes. Each mortality assumption is correct, but each includes a common shock component with constant force 0.02.

Calculate the increase in the actuarial present value over what you originally calculated.

- (A) 0.020
- (B) 0.039
- (C) 0.093
- (D) 0.109
- (E) 0.163

123. The number of accidents follows a Poisson distribution with mean 12. Each accident generates 1, 2, or 3 claimants with probabilities $\frac{1}{2}$, $\frac{1}{3}$, $\frac{1}{6}$, respectively.

Calculate the variance in the total number of claimants.

- (A) 20
- (B) 25
- (C) 30
- (D) 35
- (E) 40

124. For a claims process, you are given:

- (i) The number of claims $\{N(t), t \geq 0\}$ is a nonhomogeneous Poisson process with intensity function:

$$\lambda(t) = \begin{cases} 1, & 0 \leq t < 1 \\ 2, & 1 \leq t < 2 \\ 3, & 2 \leq t \end{cases}$$

- (ii) Claims amounts Y_i are independently and identically distributed random variables that are also independent of $N(t)$.
- (iii) Each Y_i is uniformly distributed on $[200, 800]$.
- (iv) The random variable P is the number of claims with claim amount less than 500 by time $t = 3$.
- (v) The random variable Q is the number of claims with claim amount greater than 500 by time $t = 3$.
- (vi) R is the conditional expected value of P , given $Q = 4$.

Calculate R .

- (A) 2.0
- (B) 2.5
- (C) 3.0
- (D) 3.5
- (E) 4.0

125. Lottery Life issues a special fully discrete whole life insurance on (25):

- (i) At the end of the year of death there is a random drawing. With probability 0.2, the death benefit is 1000. With probability 0.8, the death benefit is 0.
- (ii) At the start of each year, including the first, while (25) is alive, there is a random drawing. With probability 0.8, the level premium π is paid. With probability 0.2, no premium is paid.
- (iii) The random drawings are independent.
- (iv) Mortality follows the Illustrative Life Table.
- (v) $i = 0.06$
- (vi) π is determined using the equivalence principle.

Calculate the benefit reserve at the end of year 10.

- (A) 10.25
- (B) 20.50
- (C) 30.75
- (D) 41.00
- (E) 51.25

126. A government creates a fund to pay this year's lottery winners.

You are given:

- (i) There are 100 winners each age 40.
- (ii) Each winner receives payments of 10 per year for life, payable annually, beginning immediately.
- (iii) Mortality follows the Illustrative Life Table.
- (iv) The lifetimes are independent.
- (v) $i = 0.06$
- (vi) The amount of the fund is determined, using the normal approximation, such that the probability that the fund is sufficient to make all payments is 95%.

Calculate the initial amount of the fund.

- (A) 14,800
- (B) 14,900
- (C) 15,050
- (D) 15,150
- (E) 15,250

127. For a special fully discrete 35-payment whole life insurance on (30):

- (i) The death benefit is 1 for the first 20 years and is 5 thereafter.
- (ii) The initial benefit premium paid during the each of the first 20 years is one fifth of the benefit premium paid during each of the 15 subsequent years.
- (iii) Mortality follows the Illustrative Life Table.
- (iv) $i = 0.06$
- (v) $A_{30:\overline{20}|} = 0.32307$
- (vi) $\ddot{a}_{30:\overline{35}|} = 14.835$

Calculate the initial annual benefit premium.

- (A) 0.010
- (B) 0.015
- (C) 0.020
- (D) 0.025
- (F) 0.030

128. For independent lives (x) and (y) :

- (i) $q_x = 0.05$
- (ii) $q_y = 0.10$
- (iii) Deaths are uniformly distributed over each year of age.

Calculate ${}_{0.75}q_{xy}$.

- (A) 0.1088
- (B) 0.1097
- (C) 0.1106
- (D) 0.1116
- (E) 0.1125

129. In a clinic, physicians volunteer their time on a daily basis to provide care to those who are not eligible to obtain care otherwise. The number of physicians who volunteer in any day is uniformly distributed on the integers 1 through 5. The number of patients that can be served by a given physician has a Poisson distribution with mean 30.

Determine the probability that 120 or more patients can be served in a day at the clinic, using the normal approximation with continuity correction.

- (A) $1 - \Phi(0.68)$
- (B) $1 - \Phi(0.72)$
- (C) $1 - \Phi(0.93)$
- (D) $1 - \Phi(3.13)$
- (E) $1 - \Phi(3.16)$

- 130.** A person age 40 wins 10,000 in the actuarial lottery. Rather than receiving the money at once, the winner is offered the actuarially equivalent option of receiving an annual payment of K (at the beginning of each year) guaranteed for 10 years and continuing thereafter for life.

You are given:

- (i) $i = 0.04$
- (ii) $A_{40} = 0.30$
- (iii) $A_{50} = 0.35$
- (iv) $A_{40:\overline{10}|}^1 = 0.09$

Calculate K .

- (A) 538
 - (B) 541
 - (C) 545
 - (D) 548
 - (E) 551
- 131.** Mortality for Audra, age 25, follows De Moivre's law with $\omega = 100$. If she takes up hot air ballooning for the coming year, her assumed mortality will be adjusted so that for the coming year only, she will have a constant force of mortality of 0.1.

Calculate the decrease in the 11-year temporary complete life expectancy for Audra if she takes up hot air ballooning.


- (A) 0.10
- (B) 0.35
- (C) 0.60
- (D) 0.80
- (E) 1.00

132. For a 5-year fully continuous term insurance on (x):


- (i) $\delta = 0.10$
- (ii) All the graphs below are to the same scale.
- (iii) All the graphs show $\mu_x(t)$ on the vertical axis and t on the horizontal axis.

Which of the following mortality assumptions would produce the highest benefit reserve at the end of year 2?


(A)




(B)




(C)



(D)



(E)



133. For a multiple decrement table, you are given:

- (i) Decrement (1) is death, decrement (2) is disability, and decrement (3) is withdrawal.
- (ii) $q'_{60}{}^{(1)} = 0.010$
- (iii) $q'_{60}{}^{(2)} = 0.050$
- (iv) $q'_{60}{}^{(3)} = 0.100$
- (v) Withdrawals occur only at the end of the year.
- (vi) Mortality and disability are uniformly distributed over each year of age in the associated single decrement tables.

Calculate $q_{60}^{(3)}$.

- (A) 0.088
- (B) 0.091
- (C) 0.094
- (D) 0.097
- (E) 0.100

134. The number of claims, N , made on an insurance portfolio follows the following distribution:

n	$\Pr(N=n)$
0	0.7
2	0.2
3	0.1

If a claim occurs, the benefit is 0 or 10 with probability 0.8 and 0.2, respectively.

The number of claims and the benefit for each claim are independent.

Calculate the probability that aggregate benefits will exceed expected benefits by more than 2 standard deviations.

- (A) 0.02
- (B) 0.05
- (C) 0.07
- (D) 0.09
- (E) 0.12

135. For a special whole life insurance of 100,000 on (x) , you are given:

- (i) $\delta = 0.06$
- (ii) The death benefit is payable at the moment of death.
- (iii) If death occurs by accident during the first 30 years, the death benefit is doubled.
- (iv) $\mu_x^{(\tau)}(t) = 0.008, t \geq 0$
- (v) $\mu_x^{(1)}(t) = 0.001, t \geq 0$, where $\mu_x^{(1)}$ is the force of decrement due to death by accident.

Calculate the single benefit premium for this insurance.

- (A) 11,765
- (B) 12,195
- (C) 12,622
- (D) 13,044
- (E) 13,235

136. You are given the following extract from a select-and-ultimate mortality table with a 2-year select period:

x	$l_{[x]}$	$l_{[x]+1}$	l_{x+2}	$x + 2$
60	80,625	79,954	78,839	62
61	79,137	78,402	77,252	63
62	77,575	76,770	75,578	64

Assume that deaths are uniformly distributed between integral ages.

Calculate ${}_{0.9}q_{[60]+0.6}$.

- (A) 0.0102
- (B) 0.0103
- (C) 0.0104
- (D) 0.0105
- (E) 0.0106

137. A claim count distribution can be expressed as a mixed Poisson distribution. The mean of the Poisson distribution is uniformly distributed over the interval $[0,5]$.

Calculate the probability that there are 2 or more claims.

- (A) 0.61
- (B) 0.66
- (C) 0.71
- (D) 0.76
- (E) 0.81

138. For a double decrement table with $l_{40}^{(\tau)} = 2000$:

x	$q_x^{(1)}$	$q_x^{(2)}$	$q_x'^{(1)}$	$q_x'^{(2)}$
40	0.24	0.10	0.25	y
41	--	--	0.20	$2y$

Calculate $l_{42}^{(\tau)}$.

- (A) 800
- (B) 820
- (C) 840
- (D) 860
- (E) 880

139. For a fully discrete whole life insurance of 10,000 on (30):

- (i) π denotes the annual premium and $L(\pi)$ denotes the loss-at-issue random variable for this insurance.
- (ii) Mortality follows the Illustrative Life Table.
- (iii) $i=0.06$

Calculate the lowest premium, π' , such that the probability is less than 0.5 that the loss $L(\pi')$ is positive.

- (A) 34.6
- (B) 36.6
- (C) 36.8
- (D) 39.0
- (E) 39.1

140. Y is the present-value random variable for a special 3-year temporary life annuity-due on (x) . You are given:

(i) ${}_t p_x = 0.9^t, \quad t \geq 0$

(ii) K is the curtate-future-lifetime random variable for (x) .

(iii) $Y = \begin{cases} 1.00, & K = 0 \\ 1.87, & K = 1 \\ 2.72, & K = 2, 3, \dots \end{cases}$

Calculate $\text{Var}(Y)$.

(A) 0.19

(B) 0.30

(C) 0.37

(D) 0.46

(E) 0.55

141. A claim severity distribution is exponential with mean 1000. An insurance company will pay the amount of each claim in excess of a deductible of 100.

Calculate the variance of the amount paid by the insurance company for one claim, including the possibility that the amount paid is 0.

(A) 810,000

(B) 860,000

(C) 900,000

(D) 990,000

(E) 1,000,000

142. For a fully continuous whole life insurance of 1 on (x) :

- (i) π is the benefit premium.
- (ii) L is the loss-at-issue random variable with the premium equal to π .
- (iii) L^* is the loss-at-issue random variable with the premium equal to 1.25π .
- (iv) $\bar{a}_x = 5.0$
- (v) $\delta = 0.08$
- (vi) $\text{Var}(L) = 0.5625$

Calculate the sum of the expected value and the standard deviation of L^* .

- (A) 0.59
- (B) 0.71
- (C) 0.86
- (D) 0.89
- (E) 1.01

143. Workers' compensation claims are reported according to a Poisson process with mean 100 per month. The number of claims reported and the claim amounts are independently distributed. 2% of the claims exceed 30,000.

Calculate the number of complete months of data that must be gathered to have at least a 90% chance of observing at least 3 claims each exceeding 30,000.

- (A) 1
- (B) 2
- (C) 3
- (D) 4
- (E) 5

144. For students entering a three-year law school, you are given:

(i) The following double decrement table:

Academic Year	For a student at the beginning of that academic year, probability of		
	Academic Failure	Withdrawal for All Other Reasons	Survival Through Academic Year
1	0.40	0.20	--
2	--	0.30	--
3	--	--	0.60

(ii) Ten times as many students survive year 2 as fail during year 3.

(iii) The number of students who fail during year 2 is 40% of the number of students who survive year 2.

Calculate the probability that a student entering the school will withdraw for reasons other than academic failure before graduation.

- (A) Less than 0.35
- (B) At least 0.35, but less than 0.40
- (C) At least 0.40, but less than 0.45
- (D) At least 0.45, but less than 0.50
- (E) At least 0.50

145. Given:

(i) Superscripts M and N identify two forces of mortality and the curtate expectations of life calculated from them.

$$(ii) \quad \mu_{25}^N(t) = \begin{cases} \mu_{25}^M(t) + 0.1*(1-t) & 0 \leq t \leq 1 \\ \mu_{25}^M(t) & t > 1 \end{cases}$$

$$(iii) \quad e_{25}^M = 10.0$$

Calculate e_{25}^N .

(A) 9.2

(B) 9.3

(C) 9.4

(D) 9.5

(E) 9.6

146. A fund is established to pay annuities to 100 independent lives age x . Each annuitant will receive 10,000 per year continuously until death. You are given:

(i) $\delta = 0.06$

(ii) $\bar{A}_x = 0.40$

(iii) ${}^2\bar{A}_x = 0.25$

Calculate the amount (in millions) needed in the fund so that the probability, using the normal approximation, is 0.90 that the fund will be sufficient to provide the payments.

(A) 9.74

(B) 9.96

(C) 10.30

(D) 10.64

(E) 11.10

- 147.** Total hospital claims for a health plan were previously modeled by a two-parameter Pareto distribution with $\alpha = 2$ and $\theta = 500$.

The health plan begins to provide financial incentives to physicians by paying a bonus of 50% of the amount by which total hospital claims are less than 500. No bonus is paid if total claims exceed 500.

Total hospital claims for the health plan are now modeled by a new Pareto distribution with $\alpha = 2$ and $\theta = K$. The expected claims plus the expected bonus under the revised model equals expected claims under the previous model.

Calculate K .

- (A) 250
- (B) 300
- (C) 350
- (D) 400
- (E) 450

- 148.** A decreasing term life insurance on (80) pays $(20-k)$ at the end of the year of death if (80) dies in year $k+1$, for $k=0,1,2,\dots,19$.

You are given:

- (i) $i=0.06$
- (ii) For a certain mortality table with $q_{80} = 0.2$, the single benefit premium for this insurance is 13.
- (iii) For this same mortality table, except that $q_{80} = 0.1$, the single benefit premium for this insurance is P .

Calculate P .

- (A) 11.1
- (B) 11.4
- (C) 11.7
- (D) 12.0
- (E) 12.3

- 149.** Job offers for a college graduate arrive according to a Poisson process with mean 2 per month. A job offer is acceptable if the wages are at least 28,000. Wages offered are mutually independent and follow a lognormal distribution with $\mu = 10.12$ and $\sigma = 0.12$.

Calculate the probability that it will take a college graduate more than 3 months to receive an acceptable job offer.

- (A) 0.27
- (B) 0.39
- (C) 0.45
- (D) 0.58
- (E) 0.61

- 150.** For independent lives (50) and (60):

$$\mu(x) = \frac{1}{100-x}, \quad 0 \leq x < 100$$

Calculate $e_{\overline{50:60}}^{\circ}$.

- (A) 30
- (B) 31
- (C) 32
- (D) 33
- (E) 34

151. For an industry-wide study of patients admitted to hospitals for treatment of cardiovascular illness in 1998, you are given:

(i)

Duration In Days	Number of Patients Remaining Hospitalized
0	4,386,000
5	1,461,554
10	486,739
15	161,801
20	53,488
25	17,384
30	5,349
35	1,337
40	0

(ii) Discharges from the hospital are uniformly distributed between the durations shown in the table.

Calculate the mean residual time remaining hospitalized, in days, for a patient who has been hospitalized for 21 days.

- (A) 4.4
- (B) 4.9
- (C) 5.3
- (D) 5.8
- (E) 6.3

152. For an individual over 65:

- (i) The number of pharmacy claims is a Poisson random variable with mean 25.
- (ii) The amount of each pharmacy claim is uniformly distributed between 5 and 95.
- (iii) The amounts of the claims and the number of claims are mutually independent.

Determine the probability that aggregate claims for this individual will exceed 2000 using the normal approximation.

- (A) $1 - \Phi(1.33)$
- (B) $1 - \Phi(1.66)$
- (C) $1 - \Phi(2.33)$
- (D) $1 - \Phi(2.66)$
- (E) $1 - \Phi(3.33)$

153. For a fully discrete three-year endowment insurance of 10,000 on (50), you are given:

- (i) $i = 0.03$
- (ii) $1000q_{50} = 8.32$
- (iii) $1000q_{51} = 9.11$
- (iv) $10,000 {}_1V_{50:\overline{3}|} = 3209$
- (v) $10,000 {}_2V_{50:\overline{3}|} = 6539$
- (vi) ${}_0L$ is the prospective loss random variable at issue, based on the benefit premium.

Calculate the variance of ${}_0L$.

- (A) 277,000
- (B) 303,000
- (C) 357,000
- (D) 403,000
- (E) 454,000

- 154.** For a special 30-year deferred annual whole life annuity-due of 1 on (35):
- (i) If death occurs during the deferral period, the single benefit premium is refunded without interest at the end of the year of death.
 - (ii) $\ddot{a}_{65} = 9.90$
 - (iii) $A_{35:\overline{30}|} = 0.21$
 - (iv) $A_{35:\overline{30}|}^1 = 0.07$

Calculate the single benefit premium for this special deferred annuity.

- (A) 1.3
- (B) 1.4
- (C) 1.5
- (D) 1.6
- (E) 1.7

155. Given:

- (i) $\mu(x) = F + e^{2x}, \quad x \geq 0$
- (ii) ${}_{0.4}p_0 = 0.50$

Calculate F .

- (A) -0.20
- (B) -0.09
- (C) 0.00
- (D) 0.09
- (E) 0.20

156-157 Use the following information for questions 156 and 157.

An insurer has excess-of-loss reinsurance on auto insurance. You are given:

- (i) Total expected losses in the year 2001 are 10,000,000.
- (ii) In the year 2001 individual losses have a Pareto distribution with

$$F(x) = 1 - \left(\frac{2000}{x + 2000} \right)^2, \quad x > 0.$$

- (iii) Reinsurance will pay the excess of each loss over 3000.
- (iv) Each year, the reinsurer is paid a ceded premium, C_{year} , equal to 110% of the expected losses covered by the reinsurance.
- (v) Individual losses increase 5% each year due to inflation.
- (vi) The frequency distribution does not change.

156. Calculate C_{2001} .

- (A) 2,200,000
- (B) 3,300,000
- (C) 4,400,000
- (D) 5,500,000
- (E) 6,600,000

157. Calculate C_{2002} / C_{2001} .

- (A) 1.04
- (B) 1.05
- (C) 1.06
- (D) 1.07
- (E) 1.08